

15/7/03

Biological studies on radiation similar to that emitted by cell phones

Adey WR, Byus CV, Cain CD, Higgins RJ, Jones RA, Kean CJ, Kuster N, MacMurray A, Stagg RB, Zimmerman G, Phillips JL, Haggren W, Spontaneous and nitrosourea-induced primary tumors of the central nervous system in Fischer 344 rats chronically exposed to 836 MHz modulated microwaves. *Radiat Res* 152(3):293-302, 1999.

We have tested an 836.55 MHz field with North American Digital Cellular (NADC) modulation in a 2-year animal bioassay that included fetal exposure. In offspring of pregnant Fischer 344 rats, we tested both spontaneous tumorigenicity and the incidence of induced central nervous system (CNS) tumors after a single dose of the carcinogen ethylnitrosourea (ENU) in utero, followed by intermittent digital-phone field exposure for 24 months. Far-field exposures began on gestational day 19 and continued until weaning at age 21 days. Near-field exposures began at 35 days and continued for the next 22 months, 4 consecutive days weekly, 2 h/day. SAR levels simulated localized peak brain exposures of a cell phone user. Of the 236 original rats, 182 (77%) survived to the termination of the whole experiment and were sacrificed at age 709-712 days. The 54 rats (23%) that died during the study ("preterm rats") formed a separate group for some statistical analyses. There was no evidence of tumorigenic effects in the CNS from exposure to the TDMA field. However, some evidence of tumor-inhibiting effects of TDMA exposure was apparent. Overall, the TDMA field-exposed animals exhibited trends toward a reduced incidence of spontaneous CNS tumors ($P < 0.16$, two-tailed) and ENU-induced CNS tumors ($P < 0.16$, two-tailed). In preterm rats, where primary neural tumors were determined to be the cause of death, fields decreased the incidence of ENU-induced tumors ($P < 0.03$, two-tailed). We discuss a possible approach to evaluating with greater certainty the possible inhibitory effects of TDMA-field exposure on tumorigenesis in the CNS.

Adey WR, Byus CV, Cain CD, Higgins RJ, Jones RA, Kean CJ, Kuster N, MacMurray A, Stagg RB, Zimmerman G, Spontaneous and nitrosourea-induced primary tumors of the central nervous system in Fischer 344 rats exposed to frequency-modulated microwave fields. *Cancer Res* 60(7):1857-1863, 2000.

In a 2-year bioassay, we exposed Fischer 344 rats to a frequency-modulated (FM) signal (836.55 MHz \pm 12.5 KHz deviation) simulating radiofrequency exposures in the head of users of hand-held mobile phones. We tested for effects on spontaneous tumorigenicity of central nervous system (CNS) tumors in the offspring of pregnant rats and also for modified incidence of primary CNS tumors in rats treated with a single dose of the neurocarcinogen ethylnitrosourea (ENU) in utero. ENU dosage (4 mg/kg) was selected to give an expected brain

tumor incidence of 10-15% over the mean life span of 26 months. Pregnant dams (n = 102) were randomly assigned to six groups. Their offspring were treated as cohorts in each of the six groups (n = 90 per group; total, n = 540): Sham ENU/Sham Field, Sham ENU/Field Exposed, ENU/Sham Field, ENU/Field Exposed, ENU/Cage Control, and Sham ENU/Cage Control. Intermittent field exposures began on gestation day 19 and continued until weaning at 21 days, resuming thereafter at 31 days and continuing until experiment termination at 731-734 days. Energy absorption rates (SARs) in the rats' brains were similar to localized peak brain exposures of a phone user (female, 236 g, 1.0 W/kg; male, 450 g, 1.2 W/kg). Of the original 540 rats, 168 died before the termination of the experiment. In these rats, ENU significantly reduced survival from a mean of 708 days in three groups without ENU treatment to 645 days in three groups treated with ENU ($P < 0.0005$). There were no effects on survival attributable to FM field exposure in either ENU-treated or in sham-treated groups. Spontaneous CNS tumor incidence in control groups was 1.1-4.4% but sharply higher in rats receiving ENU (14.4-22.2%; $P < 0.0001$). No FM field-mediated changes were observed in number, incidence, or histological type of either spontaneous or ENU-induced brain tumors, nor were gender differences detected in tumor numbers. These negative findings with FM fields contrast with our study using standard digital phone fields pulsed on and off at 50/se, where a trend was noted toward reduced incidence of both spontaneous and ENU-induced CNS tumors (W. R. Adey et al., *Radiat. Res.*, 152: 293-302, 1999). Although consistent but not attaining significance in the experiment overall (spontaneous CNS tumors, $P < 0.08$ one-tailed; $P < 0.16$ two-tailed; ENU-induced CNS tumors, $P < 0.08$ one-tailed, $P < 0.16$ two-tailed), the trend was significant ($P < 0.015$ one-tailed, $P < 0.03$, two-tailed) in rats that received ENU and died prior to experiment termination, with a primary brain tumor as the cause of death. We discuss differences in the signaling structure of digital and FM fields. Certain bioeffects induced by either amplitude-modulated or pulsed radiofrequency fields at athermal levels have not been seen with fields of similar average power but unvarying in intensity (continuous wave or frequency-modulated fields).

Anane R, Geffard M, Taxile M, Bodet D, Billaudel B, Dulou PE, Veyret B. Effects of GSM-900 microwaves on the experimental allergic encephalomyelitis (EAE) rat model of multiple sclerosis.

***Bioelectromagnetics* 24(3):211-213, 2003.** The effects of acute exposure to GSM-900 microwaves (900 MHz, 217 Hz pulse modulation) on the clinical parameters of the acute experimental allergic encephalomyelitis (EAE) model in rats were investigated in two independent experiments: rats were either habituated or nonhabituated to the exposure restrainers. EAE was induced with a mixture of myelin basic protein and Mycobacterium tuberculosis. Female Lewis rats were divided into cage control, sham exposed, and two groups exposed either at 1.5 or 6.0 W/kg local specific absorption rate (SAR averaged over the brain) using a loop antenna placed over their heads. There was no effect of a 21-day exposure (2 h/day) on the onset, duration, and termination of the EAE crisis.

Antonopoulos A, Eisenbrandt H, Obe G, Effects of high-frequency electromagnetic fields on human lymphocytes in vitro. *Mutat Res* 395(2-3): 209-214, 1997.

Human peripheral lymphocytes were incubated in the presence of high-frequency electromagnetic fields of 380, 900 and 1800 MHz. The measured endpoints were cell cycle progression and the frequencies of sister-chromatid exchanges. No differences between treated and control cultures could be found.

Auvinen A, Hietanen M, Luukkonen R, Koskela R-S, Brain Tumors and Salivary Gland Cancers Among Cellular Telephone Users *Epidemiology* 13:356-359, 2002.

Background. Possible risk of cancer associated with use of cellular telephones has lately been a subject of public debate. Methods. We conducted a register-based, case-control study on cellular phone use and cancer. The study subjects were all cases of brain tumor (N = 398) and salivary gland cancer (N = 34) diagnosed in Finland in 1996, with five controls per case. Results. Cellular phone use was not associated with brain tumors or salivary gland cancers overall, but there was a weak association between gliomas and analog cellular phones. Conclusions. A register-based approach has limited value in risk assessment of cellular phone use owing to lack of information on exposure.

Bartsch H, Bartsch C, Seebald E, Deerberg F, Dietz K, Vollrath L, Mecke D. Chronic Exposure to a GSM-like Signal (Mobile Phone) Does Not Stimulate the Development of DMBA-Induced Mammary Tumors in Rats: Results of Three Consecutive Studies. *Radiat Res* 157(2):183-190, 2002.

Certain epidemiological and experimental studies raised concerns about the safety of radiofrequency (RF) electromagnetic fields because of a possible increased risk of leukemia and lymphoma. In this study, an RF field used in mobile telecommunication was tested using 7,12-dimethylbenz[a]anthracene (DMBA)-induced mammary tumors in female Sprague-Dawley rats as a model for human breast cancer. Three experiments were carried out under strictly standardized conditions and were started on the same day of three consecutive years. The field consisted of a GSM-like signal (900 MHz pulsed at 217 Hz, pulse width 577 [μ s]) of relatively low power flux density (100 [μ W/cm²] [plus minus] 3 dB) and was applied continuously throughout each experiment to freely moving animals. The specific absorption rates averaged over the whole body were 17.5--70 mW/kg. The highest values in young animals were at or around the exposure limit permissible for the general public (i.e. 80 mW/kg). The animals were palpated weekly for the presence of mammary tumors and were killed humanely when tumors reached a diameter of 1--2 cm to allow a reliable histopathological classification and a distinction between malignant and benign subtypes. The overall results of the three studies are that there was no statistically significant effect of RF-field exposure on tumor latency and that the cumulative tumor incidence at the end of the experiment was unaffected as well. The risk ratios were 1.08 (95% CI: 0.91--1.29) and 0.96 (95% CI: 0.85--1.07) for benign and malignant tumors, respectively. These observations are in agreement with other published findings. In the first experiment, however, the median latency for the development of the first malignant tumor in each animal was statistically significantly extended for RF-field-exposed animals compared to controls (278 days compared to 145 days, P = 0.009). No such differences were detected in the two subsequent experiments. These results show that low-level RF radiation does not appear to possess carcinogenic or cancer-promoting effects on DMBA-induced mammary tumors. To explain the mechanisms underlying the different results obtained in the three experiments, a hypothesis is presented which is based upon the neuroendocrine control mechanisms involved in the promotion of DMBA-induced mammary tumors. Despite the apparent absence of stimulatory effects of low-level RF-field exposure on the development and growth of solid tumors, it will be necessary to verify these results for leukemias and lymphomas, which may have completely different biological control mechanisms.

Beason RC, Semm P. Responses of neurons to an amplitude modulated microwave stimulus. *Neurosci Lett* 333(3):175-178, 2002. In this study we investigated the effects of a pulsed radio frequency signal similar to the signal produced by global system for mobile communication telephones (900 MHz carrier, modulated at 217 Hz) on neurons of the avian brain. We found that such stimulation resulted in changes in the amount of neural activity by more than half of the brain cells. Most (76%) of the responding cells increased their rates of firing by an average 3.5-fold. The other responding cells exhibited a decrease in their rates of spontaneous activity. Such responses indicate potential effects on humans using hand-held cellular phones.

Bisht KS, Moros EG, Straube WL, Baty JD, Roti Roti JL, The Effect of 835.62 MHz FDMA or 847.74 MHz CDMA Modulated Radiofrequency Radiation on the Induction of Micronuclei in C3H 10T_{1/2} Cells. *Radiat. Res.* 157, 506–515, 2002.

To determine if radiofrequency (RF) radiation induces the formation of micronuclei, C3H 10T_{1/2} cells were exposed to 835.62 MHz frequency division multiple access (FDMA) or 847.74 MHz code division multiple access (CDMA) modulated RF radiation. After the exposure to RF radiation, the micronucleus assay was performed by the cytokinesis block method using cytochalasin B treatment. The micronuclei appearing after mitosis were scored in binucleated cells using acridine orange staining. The frequency of micronuclei was scored both as the percentage of binucleated cells with micronuclei and as the number of micronuclei per 100 binucleated cells. Treatment of cells with cytochalasin B at a concentration of 2 μ g/ml for 22 h was found to yield the maximum number of binucleated cells in C3H 10T_{1/2} cells. The method used for the micronucleus assay in the present study detected a highly significant dose response for both indices of micronucleus production in the dose range of 0.1–1.2 Gy and it was sensitive enough to detect a significant ($P > 0.05$) increase in micronuclei after doses of 0.3 Gy in exponentially growing cells and after 0.9 Gy in plateau-phase cells. Exponentially growing cells or plateau-phase cells were exposed to CDMA (3.2 or 4.8 W/kg) or FDMA (3.2 or 5.1 W/kg) RF radiation for 3, 8, 16 or 24 h. In three repeat experiments, no exposure condition was found by analysis of variance to result in a significant increase relative to sham-exposed cells either in the percentage of binucleated cells with micronuclei or in the number of micronuclei per 100 binucleated cells. In this study, data from cells exposed to different RF signals at two SARs were compared to a common sham-exposed sample. We used the Dunnett's test, which is specifically designed for this purpose, and found no significant exposure-related differences for either plateau-phase cells or exponentially growing cells. Thus the results of this study are not consistent with the possibility that these RF radiations induce micronuclei.

Bolshakov MA, Alekseev SI, Bursting responses of Lymnea neurons to microwave radiation. *Bioelectromagnetics* 13(2):119-129, 1992. Microelectrode and voltage-clamp techniques were modified to record spontaneous electrical activity and ionic currents of Lymnea stagnalis neurons during exposure to a 900-MHz field in a waveguide-based apparatus. The field was pulse-modulated at repetition rates ranging from 0.5 to 110 pps, or it was applied as a continuous wave (CW). When subjected to pulsed waves (PW), rapid, burst-like changes in the firing rate of neurons occurred at SARs of a few W/kg. If the burst-like irregularity was present in the firing rate under control conditions, irradiation enhanced its probability of occurrence. The effect was dependent on modulation, but not on modulation frequency, and it had a threshold SAR near 0.5 W/kg. CW radiation had no effect on the firing rate pattern at the same SAR. Mediator-induced, current activation of acetylcholine, dopamine, serotonin, or gamma-aminobutyric-acid receptors of the neuronal soma was not altered during CW or PW exposures and, hence, could not have been responsible for the bursting effect.

Borbely, AA, Huber, R, Graf, T, Fuchs, B, Gallmann, E, Achermann, P, Pulsed high-frequency electromagnetic field affects human sleep and sleep electroencephalogram. *Neurosci Lett* 275(3):207-210, 1999.

To investigate whether the electromagnetic field (EMF) emitted by digital radiotelephone handsets affects the brain, healthy, young subjects were exposed during an entire night-time sleep episode to an intermittent radiation schedule (900 MHz; maximum specific absorption rate 1 W/kg) consisting of alternating 15-min on-15-min off intervals. Compared with a control night with sham exposure, the amount of waking after sleep onset was reduced from 18 to 12 min. Spectral power of the electroencephalogram in non-rapid eye movement sleep was increased. The maximum rise occurred in the 10-11 Hz and 13.5-14 Hz bands during the initial part of sleep and then subsided. The results demonstrate that pulsed high-frequency EMF in the range of radiotelephones may promote sleep and modify the sleep EEG.

Bornhausen M, Scheingraber H, Prenatal exposure to 900 MHz, cell-phone electromagnetic fields had no effect on operant-behavior performances of adult rats. *Bioelectromagnetics* 21(8):566-574, 2000. To clarify potential health risks of radio-frequency electromagnetic fields (EMFs) used in cellular telephone technology to the developing brain, Wistar rats were continuously exposed during pregnancy to a low-level (0.1 mW/cm²) 900 MHz, 217 Hz pulse modulated EMF that approximated the highest legal exposure of normal populations to the radiation of base antennas of the GSM digital cell-phone technology. Whole body average specific absorption rate (SAR) values for the freely roaming, pregnant animals were measured in models; they ranged between 17.5 and 75 mW/kg. The offspring of exposed and of sham-exposed dams were coded and tested later as adults in a battery of ten simultaneously operated test chambers (Skinner boxes) during night time. Eight groups of ten coded animals in each group were tested for learning deficits in a sequence of nine, computer-controlled, 15 h sessions of the food-reinforced contingency Differential Reinforcement of Rate with increasing performance requirements. Two different sets of events were recorded: The food-reinforced lever-pressing activity of the animals and the inter-response intervals (IRIs) between consecutive lever presses. IRI-occurrence patterns discriminated consistently between "learners" and "non-learners". Analyses of performance scores and of IRI-patterns both showed that exposure in-utero to the GSM field did not induce any measurable cognitive deficits.

Bortkiewicz A, Pilacik B, Gadzicka E, Szymczak W. The excretion of 6-hydroxymelatonin sulfate in healthy young men exposed to electromagnetic fields emitted by cellular phone -- an experimental study. *Neuroendocrinol Lett* 23 Suppl 1:88-91, 2002. OBJECTIVES: It is quite likely that non-visible electromagnetic fields (EMF) may affect melatonin production. Some studies confirmed this hypothesis and showed that extremely low EMF altered pineal function in animals and humans. Thus, it is reasonable to suppose

that EMF emitted by cellular phones may also influence secretion of melatonin. The present study sought to evaluate possible effect of the exposure to EMF emitted by cellular phone on 6-hydroxymelatonin sulfate (6-OHMS) excretion, which reflects melatonin levels in blood. MATERIAL AND METHODS: The examined group consisted of 9 healthy males aged 19-29 years. The experiment was performed under controlled conditions (the light intensity-50 lx till midnight and 0 lx during night). Each person was examined twice: on a day without exposure (control day, C-day) and on a day with continuous exposure (60 min. exposure from cellular phone, frequency 900 MHz, pulsed with 217 Hz, pulse with 576 micros, SAR 1.23 W/kg, E-day). From 7 p.m. to 8 p.m. they used a cellular phone. The subjects did not know which day was E-day, and which was C-day. From 8 p.m. till midnight the subjects listened to music and then they slept till 7 a.m. next day. Urine samples were collected at 7 p.m., at midnight, and at 7 a.m. in the same way in C-day as in E-day. Samples were frozen for later ELISA analysis of 6-OHMS. The 6-OHMS ELISA kit from Immuno-Biological Laboratories (Hamburg) was used for measurement of 6-OHMS. The data were analysed using Wilcoxon matched-pairs signed-ranks test for each subject and for the whole group. We compared 6-OHMS level on the E-day and on the C-day separately for 3 time-points - 7 p.m., midnight, 7 a.m. RESULTS: Mean 6-OHMS level in both experiments did not differ significantly for any of the respective time points. Circadian variations of 6-OHMS level were detected in all subjects. CONCLUSIONS: The results of our investigation have demonstrated that EMF emitted by cellular phones has no distinct influence on the melatonin level.

Braune, S, Wrocklage, C, Raczek, J, Gailus, T, Lucking, CH, Resting blood pressure increase during exposure to a radio-frequency electromagnetic field. *Lancet* 351(9119):1857-1858, 1998.

Exposure of the right hemisphere to a radiofrequency EMF for 35 min causes in human subjects an increase in sympathetic efferent activity with increases in the resting blood pressure between 5-10 mm Hg. The effect is likely caused by vasoconstriction.

Braune S, Riedel A, Schulte-Monting J, Raczek J. Influence of a radiofrequency electromagnetic field on cardiovascular and hormonal parameters of the autonomic nervous system in healthy individuals. *Radiat Res* 158(3):352-356, 2002. The potential health risks of radiofrequency electromagnetic fields (EMFs) emitted by mobile phones are of considerable public interest. The present study investigated the hypothesis, based on the results of our previous study, that exposure to EMFs can increase sympathetic vasoconstrictor activity. Forty healthy young males and females underwent a single-blind, placebo-controlled protocol once on each of two different days. Each investigation included successive periods of placebo and EMF exposure, given in a randomized order. The exposure was implemented by a GSM-like signal (900 MHz, pulsed with 217 Hz, 2 W) using a mobile phone mounted on the right-hand side of the head in a typical telephoning position. Each period of placebo

exposure and of EMF exposure consisted of 20 min of supine rest, 10 min of 70 degrees upright tilt on a tilt table, and another 20 min of supine rest. Blood pressure, heart rate and cutaneous capillary perfusion were measured continuously. In addition, serum levels of norepinephrine, epinephrine, cortisol and endothelin were analyzed in venous blood samples taken every 10 min. Similar to the previous study, systolic and diastolic blood pressure each showed slow, continuous, statistically significant increases of about 5 mmHg during the course of the protocol. All other parameters either decreased in parallel or remained constant. However, analysis of variance showed that the changes in blood pressure and in all other parameters were independent of the EMF exposure. These findings do not support the assumption of a nonthermal influence of EMFs emitted by mobile phones on the cardiovascular autonomic nervous system in healthy humans.

Burch JB, Reif JS, Noonan CW, Ichinose T, Bachand AM, Koleber TL, Yost MG. Melatonin metabolite excretion among cellular telephone users. *Int J Rad Biol* 78: 1029-1036, 2002.

Abstract: *Purpose:* The relationship between cellular telephone use and excretion of the melatonin metabolite 6-hydroxymelatonin sulfate (6-OHMS) was evaluated in two populations of male electric utility workers (Study 1, $n=149$; Study 2, $n=77$). *Materials and methods:* Participants collected urine samples and recorded cellular telephone use over 3 consecutive workdays. Personal 60-Hz magnetic field (MF) and ambient light exposures were characterized on the same days using EMDEX II meters. A repeated measures analysis was used to assess the effects of cellular telephone use, alone and combined with MF exposures, after adjustment for age, participation month and light exposure. *Results:* No change in 6-OHMS excretion was observed among those with daily cellular telephone use >25 min in Study 1 (5 worker-days). Study 2 workers with >25 min cellular telephone use per day (13 worker-days) had lower creatinine-adjusted mean nocturnal 6-OHMS concentrations ($p=0.05$) and overnight 6-OHMS excretion ($p=0.03$) compared with those without cellular telephone use. There was also a linear trend of decreasing mean nocturnal 6-OHMS/creatinine concentrations ($p=0.02$) and overnight 6-OHMS excretion ($p=0.08$) across categories of increasing cellular telephone use. A combined effect of cellular telephone use and occupational 60-Hz MF exposure in reducing 6-OHMS excretion was also observed in Study 2. *Conclusions:* Exposure-related reductions in 6-OHMS excretion were observed in Study 2, where daily cellular telephone use of >25 min was more prevalent. Prolonged use of cellular telephones may lead to reduced melatonin production, and elevated 60-Hz MF exposures may potentiate the effect.

Cain CD, Thomas DL, Adey WR, Focus formation of C3H/10T1/2 cells and exposure to a 836.55 MHz modulated radiofrequency field. *Bioelectromagnetics* 18(3):237-243, 1997.

Disruption of communication between transformed cells and normal cells is involved in tumor promotion. We have tested the hypothesis that exposures

to radiofrequency (RF) fields using a form of digital modulation (TDMA) and a chemical tumor promoter, 12-O-tetradecanoylphorbol-13-acetate (TPA), are copromoters that enhance focus formation of transformed cells in coculture with parental C3H/10T1/2 murine fibroblasts. RF field exposures did not influence TPA's dose-dependent promotion of focus formation in coculture. Cell cultures were exposed to an 836.55 MHz TDMA-modulated field in TEM transmission line chambers, with incident energies that simulated field intensities at a user's head. Specific absorption rates (SARs) of 0.15, 1.5, and 15 $\mu\text{W/g}$ were used during each digital packet, and the packet frequency was 50/s. The TEM chambers were placed in a commercial incubator at 37 degrees C and 95% humidity/5% CO_2 . The RF field exposures were in a repeating cycle, 20 min on, 20 min off, 24 h/day for 28 days. At 1.5 $\mu\text{W/g}$, TPA-induced focus formation (at 10, 30, and 50 ng/ml) was not significantly different in RF-exposed cultures compared to parallel sham-exposed cultures in ten independent experiments in terms of the number, density, and area of foci. Similarly, at 0.15 and 15.0 $\mu\text{W/g}$, in two and four experiments, respectively, RF exposure did not alter TPA-induced focus formation. The findings support a conclusion that repeated exposures to this RF field do not influence tumor promotion in vitro, based on the RF field's inability to enhance TPA-induced focus formation.

Chagnaud JL, Veyret B In vivo exposure of rats to GSM-modulated microwaves: flow cytometry analysis of lymphocyte subpopulations and of mitogen stimulation. *Int J Radiat Biol* 75(1):111-113, 1999.

The effects of GSM-modulated microwaves on lymphocyte sub-populations of Sprague-Dawley rats and their normal mitogenic responses were investigated using flow cytometry analysis and a colorimetric method. No alterations were found in the surface phenotype of splenic lymphocytes or in their mitogenic activity, indicating that low-level pulsed microwaves do not seem to affect the integrity of the immune system.

Chagnaud, JL, Moreau, JM, Veyret, B, No effect of short-term exposure to GSM-modulated low-power microwaves on benzo(a)pyrene-induced tumours in rat. *Int J Radiat Biol* 75(10):1251-1256, 1999.

PURPOSE: In view of current interest in the biological effects of amplitude-modulated microwaves arising from the rapid development of mobile communications, the effects of low-level microwaves on cancer development were investigated using a rat sarcoma model. **MATERIALS AND METHODS:** Two-month-old female Sprague-Dawley rats were treated by injection of benzo(a)pyrene and irradiated with GSM (Global System for Mobile)-modulated 900-MHz microwaves in an anechoic chamber at 55 or 200 $\mu\text{W cm}^{-2}$ (75 and 270 mW kg^{-1}) average whole-body SAR, 2h daily for 2 weeks). Rats were exposed from day 20, 40 or 75 after carcinogen injection. Additional groups of rats were sham-exposed in a second anechoic chamber. Anti-phosphatidylinositol autoantibody levels were evaluated in sera to monitor malignant transformation. **RESULTS:** Microwave exposure had no effect on the development of tumours. No acceleration or delays in tumour onset were observed. Animal survival was not modified and serum autoantibody levels were similar in exposed and sham-exposed groups.

CONCLUSION: Low-level GSM microwave exposure of rat bearing benzo(a)pyrene-induced tumours had no effect on auto-antibody levels, tumour appearance and survival. The low exposure levels used here correspond to exposure limits for whole-body exposure of humans.

Chia SE, Chia HP, Tan JS, Prevalence of headache among handheld cellular telephone users in singapore: A community study. *Environ Health Perspect* 108(11):1059-1062, 2000. We carried out a cross-sectional community study in Singapore to determine the prevalence of specific central nervous system (CNS) symptoms among hand-held cellular telephone (HP) users compared to nonusers and to study the association of risk factors and CNS symptoms among HP users. A total of 808 men and women between 12 and 70 years of age, who lived in one community, were selected using one-stage cluster random sampling and responses to a structured questionnaire. The prevalence of HP users was 44.8%. Headache was the most prevalent symptom among HP users compared to non-HP users, with an adjusted prevalence rate ratio of 1.31 [95% confidence interval, 1.00-1.70]. There is a significant increase in the prevalence of headache with increasing duration of usage (in minutes per day). Prevalence of headache was reduced by more than 20% among those who used hand-free equipment for their cellular telephones as compared to those who never use the equipment. The use of HPs is not associated with a significant increase of CNS symptoms other than headache.

Cook A, Woodward A, Pearce N, Marshall C. Cellular telephone use and time trends for brain, head and neck tumours. *N Z Med J*. 116(1175):U457, 2003.

AIM: The objective of this study was to determine whether incidence rates of head and neck malignancies in New Zealand have varied since the introduction of cellular telephones in 1987. In particular, we sought to compare trends in tumour rates in anatomical sites that receive high, medium and low levels of cellular telephone radiation (based on dosimetry data). METHODS: We investigated whether trends in tumour incidence rates in New Zealand have varied since the introduction of cellular telephones in 1987. The exposure measure used was the proportion of cellular telephone subscribers within the national population, calculated using the number of subscribers over the study period. RESULTS: The graphs for high, medium and low exposure sites did not display any significant changes in trend patterns for either gender over the years 1986 to 1998. CONCLUSIONS: Incidence rates for malignancies arising in the head and neck, including those sites that hypothetically receive the highest levels of radio frequency radiation during cellular telephone use, have not changed materially since the introduction of cellular telephones to New Zealand. However, ecological studies of this nature are limited in many ways and a stronger study design is clearly needed to establish more exactly any elevation in risk.

Cranfield CG, Wood AW, Anderson V, Menezes KG. Effects of mobile phone type signals on calcium levels within human leukaemic T-cells (Jurkat cells). *Int J Radiat Biol* 77(12):1207-1217, 2001. PURPOSE: To test whether exposure to simulated GSM mobile phone signals (915 MHz, 2 W kg⁻¹)

influences the concentration of calcium or calcium signalling patterns in a human lymphocyte cell line. **MATERIALS AND METHODS:** The radiofrequency (RF) energy was delivered via a coaxial applicator to a perfused chamber where cells adherent to a thin glass coverslip were imaged by laser scanning confocal microscopy. Cell calcium concentration, estimated from Fluo-3 fluorescence, was monitored over two 10-min periods; control followed by exposed/sham, with exposure status assigned in a blind and randomized fashion. Both continuous wave (CW) and pulsed wave (PW) RF (on both phytohaemagglutinin-activated and unactivated cells) were studied (with an equal number of sham exposures) on 100 cells per category (total 800 cells). **RESULTS:** No significant changes were noted for the following: regression slope of calcium fluorescence; mean calcium concentration; number of calcium 'spikes' in each 10 min; or mean height of these 'spikes'. The average frequency from Fourier spectra of these periods showed significant alteration in one category only: PW exposure of activated cells. **CONCLUSIONS:** There is no clear indication that RF emissions from mobile phones are associated with any changes in calcium levels or calcium signalling in lymphocytes.

Croft R, Chandler J, Burgess A, Barry R, Williams J, Clarke A. Acute mobile phone operation affects neural function in humans. *Clin Neurophysiol* 113(10):1623, 2002. **OBJECTIVES:** Mobile phones (MP) are used extensively and yet little is known about the effects they may have on human physiology. There have been conflicting reports regarding the relation between MP use and the electroencephalogram (EEG). The present study suggests that this conflict may be due to methodological differences such as exposure durations, and tests whether exposure to an active MP affects EEG as a function of time. **METHODS:** Twenty-four subjects participated in a single-blind fully counterbalanced cross-over design, where both resting EEG and phase-locked neural responses to auditory stimuli were measured while a MP was either operating or turned off. **RESULTS:** MP exposure altered resting EEG, decreasing 1-4Hz activity (right hemisphere sites), and increasing 8-12Hz activity as a function of exposure duration (midline posterior sites). MP exposure also altered early phase-locked neural responses, attenuating the normal response decrement over time in the 4-8Hz band, decreasing the response in the 12-30Hz band globally and as a function of time, and increasing midline frontal and lateral posterior responses in the 30-45Hz band. **CONCLUSIONS:** Active MPs affect neural function in humans and do so as a function of exposure duration. The temporal nature of this effect may contribute to the lack of consistent results reported in the literature.

d'Ambrosio G, Massa R, Scarfi MR, Zeni O, Cytogenetic damage in human lymphocytes following GMSK phase modulated microwave exposure. *Bioelectromagnetics* 23:7-13, 2002.

The present study investigated, using in vitro experiments on human lymphocytes, whether exposure to a microwave frequency used for mobile communication, either unmodulated or in presence of phase only modulation, can cause modification of cell proliferation kinetics and/or genotoxic effects, by

evaluating the cytokinesis block proliferation index and the micronucleus frequency. In the GSM 1800 mobile communication systems the field is both phase (Gaussian minimum shift keying, GMSK) and amplitude (time domain multiple access, TDMA) modulated. The present study investigated only the effects of phase modulation, and no amplitude modulation was applied. Human peripheral blood cultures were exposed to 1.748 GHz, either continuous wave (CW) or phase only modulated wave (GMSK), for 15 min. The maximum specific absorption rate (~ 5 W/kg) was higher than that occurring in the head of mobile phone users; however, no changes were found in cell proliferation kinetics after exposure to either CW or GMSK fields. As far as genotoxicity is concerned, the micronucleus frequency result was not affected by CW exposure; however, a statistically significant micronucleus effect was found following exposure to phase modulated field. These results would suggest a genotoxic power of the phase modulation per se.

Dasdag, S, Ketani, MA, Akdag, Z, Ersay, AR, Sar, i I, Demirtas ,OC, Celik, MS, Whole-body microwave exposure emitted by cellular phones and testicular function of rats. *Urol Res* 27(3):219-223, 1999.

This study investigated whether there are adverse effects due to microwave exposure emitted by cellular phones in male rats. Eighteen Wistar Albino rats were separated into three groups, a sham group and two experimental groups. The rats were confined in Plexiglas cages and cellular phones were placed 0.5 cm under the cages. In the first experimental group, cellular phones were in standby position for 2 h. In the second experimental group, phones were turned to the speech position three times each for 1 min duration over 2 h. Rats in the first and second experimental groups were exposed to microwaves emitted by phones for 2 h/day for a duration of 1 month. After the last exposure the rats were killed. Brain, eyes, ears, liver, heart, lungs, stomach, kidneys, testes, small and large intestines and skin of the rats were observed histologically. The decrease of epididymal sperm counts in the speech groups were not found to be significant ($P > 0.05$). Differences in terms of normal and abnormal sperm forms were not observed ($P > 0.05$). Histological changes were especially observed in the testes of rats of the speech groups. Seminiferous tubular diameter of rat testes in the standby and speech groups was found to be lower than the sham group ($P < 0.05$). Rectal temperatures of rats in the speech group were found to be higher than the sham and standby groups ($P < 0.05$). The rectal temperatures of rats before and after exposure were also found to be significantly higher in the speech group ($P < 0.05$). Specific absorption rate (SAR) was determined as 0.141 W/kg.

Dasdag S, Zulkuf Akdag M, Aksen F, Yilmaz F, Bashan M, Mutlu Dasdag M, Salih Celik M. Whole body exposure of rats to microwaves emitted from a cell phone does not affect the testes. *Bioelectromagnetics* 24(3):182-188, 2003. The objective of this study was to investigate the effects of radiofrequency radiation emitted from cellular phones on the lipid composition, malondialdehyde

concentration, p53 immune reactivity, sperm count, morphology, histological structure of testes, and on rectal temperature of rats exposed to microwave radiation emitted from cellular phones. Sixteen Sprague-Dawley rats were separated into two groups of eight, sham exposed (control) and experimental. The rats were confined in plexiglas cages specially designed for this study, and cellular phones were placed 0.5 cm under the cages. For the experimental group, cellular phones were activated 20 min per day (7 days a week) for 1 month. For the control group, the cellular phones were placed beneath the cages for 20 min a day, but the phones were turned off. Rectal temperatures were measured weekly. For 250 mW radiated power, the whole body average SAR (rms) is 0.52 W/kg and 1 g averaged peak SAR (rms) is 3.13 W/kg. The Mann-Whitney U-test was used for statistical comparisons of groups. No statistically significant alteration in any of the endpoints was noted. This study found no evidence suggesting an adverse effect of cell phone exposure on measures of testicular function or structure.

de Seze R, Ayoub J, Peray P, Miro L, Touitou Y, Evaluation in humans of the effects of radiocellular telephones on the circadian patterns of melatonin secretion, a chronobiological rhythm marker. *J Pineal Res* 27(4):237-242, 1999.

A decrease in melatonin secretion has been observed in small mammals under exposure to extremely low frequency electromagnetic fields. As there is some concern about possible health effects of the increasing use of radiocellular telephones emitting radiofrequency electromagnetic fields, we examined whether such fields would alter melatonin levels in the human. Volunteers were two groups totalling 38 men, 20-32 yr old. Exposures were to commercially available cellular telephones of the GSM 900 type (Global System for Mobile communication at 900 MHz) or DCS 1800 type (Digital Communication System at 1800 MHz), for 2 hr/day, 5 days/wk, for 4 wk, at their maximum power. Attention of the volunteers was sustained by TV projection of movies. Blood samples were collected hourly during the night and every 3 hr in the daytime. Four sampling sessions were performed at 15-day intervals: before the beginning of the exposure period, at the middle and the end of the exposure period, and 15 days later to evaluate the persistence or late appearance of potential effects. Evaluated parameters were the maximum serum concentration, the time of this maximum, and the area under the curve of the hormone profile. Melatonin circadian profile was not disrupted in 37 young male volunteers submitted to a typical pattern of exposure to the electromagnetic fields generated by two common types of cell phones.

de Seze R, Fabbro-Peray P, Miro L, GSM radiocellular telephones do not disturb the secretion of antepituitary hormones in humans. *Bioelectromagnetics* 19(5):271-278, 1998.

It is known that the endocrine system of experimental animals is susceptible to perturbation by radiofrequency (RF) radiation. Because of the recent interest in health and safety issues of cellular telephones, an experiment was designed to evaluate the effect of a 900 MHz RF radiation emitted by a Global System for Mobile radiotelephone (217 Hz impulses, one-eighth duty cycle, 2 W peak power) on human endocrine functions. Twenty healthy male volunteers aged from 19 to 40 were inducted in the present experiment. Each subject was exposed to RF radiation through the use of a cellular phone 2 h/day, 5 days/wk, for 1 month. Subjects were their own control. End points were serum adrenocorticotropin, thyrotropin, growth hormone, prolactin, luteinizing hormone, and follicle stimulating hormone concentrations. These end points were determined in nine weekly blood samples obtained starting 3 weeks before the commencement of the exposure and ending 2 weeks after exposures. All but one blood sample was drawn 48 h after each weekly session. The seventh drawing was performed the morning after the last weekly exposure. Within each individual, the preexposure hormone concentration was used as a control. Results indicated that all hormone concentrations remained within normal physiologic ranges. A difference was not noted among the nine weekly samples in five of six hormones studied. There was a significant change only in thyrotropin concentration, showing a 21% decrease on the seventh sampling. Because this change recovered fully during the postexposure period, it is concluded that 1 month of intermittent exposures to RF radiation from a cellular telephone does not induce a long-lasting or cumulative effect on the hormone secretion rate of the anterior pituitary gland in humans.

Di Carlo A, White N, Guo F, Garrett P, Litovitz T. Chronic electromagnetic field exposure decreases HSP70 levels and lowers cytoprotection. *J. Cell. Biochem.* 84: 447-454, 2002.

Electromagnetic field (EMF) exposures have been shown to induce heat shock proteins (HSPs), which help to maintain the conformation of cellular proteins during periods of stress. We have previously reported that short-term exposure of chick embryos to either 60 Hz (extremely low frequency: ELF), or radio-frequency (RF: 915 MHz) EMFs induce protection against hypoxia. Experiments presented in the current report are based on a study in which long-term (4 days), continuous exposure to ELF-EMFs *decreased* protection against ultraviolet radiation. Based on this result, it was hypothesized that de-protection against hypoxia should also occur following long-term, continuous, or daily, repeated exposures to EMFs. To test this hypothesis, chick embryos were exposed to ELF-EMFs (8 μ T) continuously for 4 days, or to ELF or RF (3.5 mW incident power)- EMFs repeated daily (20, 30, or 60 min once or twice daily for 4 days). Several of the exposure protocols yielded embryos that had statistically significant decreases in protection against hypoxic stress (continuous and 30 or 60 min ELF twice daily; or 30 or 60 min once daily RF). This is consistent with our finding that following 4 days of ELF-EMF exposure, HSP70 levels decline by 27% as compared to controls. In addition, the superposition of ELF-EM noise,

previously shown to minimize ELF-EMF induced hypoxia protection, inhibited hypoxia de-protection caused by long term, continuous ELF or daily, repeated RF exposures. This EMF-induced decrease in HSP70 levels and resulting decline in cytoprotection suggests a mechanism by which daily exposure (such as might be experienced by mobile phone users) could enhance the probability of cancer and other diseases.

Donnellan M, McKenzie DR, French PW, Effects of exposure to electromagnetic radiation at 835 MHz on growth, morphology and secretory characteristics of a mast cell analogue, RBL-2H3. *Cell Biol Int* 21:427-439, 1997.

A mast cell line, RBL-2H3, was exposed to 835 MHz for 20 minutes, three times per day for 7 days at a power density of 8.1 ± 3 mW/cm². From day 4 onwards, it was observed that the rate of DNA synthesis and cell replication increased, that actin distribution and cell morphology became altered, and the amount of beta-hexosaminidase (a marker of granule secretion) released in response to a calcium ionophore was significantly enhanced, in comparison to unexposed cultures. There were no effects seen on levels of cytoskeletal protein synthesis or of beta-actin mRNA. Morphological changes persisted following subculture for at least 7 days in the absence of further exposure. It is hypothesized that effects of exposure to an electromagnetic field at 835 MHz may be mediated via a signal transduction pathway.

Dubreuil D, Jay T, Edeline JM. Does head-only exposure to GSM-900 electromagnetic fields affect the performance of rats in spatial learning tasks? *Behav Brain Res* 129(1-2):203-210, 2002. The rapid expansion of mobile communication has generated intense interest, but has also fuelled ongoing concerns. In both humans and animals, radiofrequency radiations are suspected to affect cognitive functions. More specifically, several studies performed in rodents have suggested that spatial learning can be impaired by electromagnetic field exposure. However, none of these previous studies have simulated the common conditions of GSM mobile phones use. This study is the first using a head-only exposure system emitting a 900-MHz GSM electromagnetic field (pulsed at 217 Hz). The two behavioural tasks that were evaluated here have been used previously to demonstrate performance deficits in spatial learning after electromagnetic field exposure: a classical radial maze elimination task and a spatial navigation task in an open-field arena (dry-land version of the Morris water maze). The performances of rats exposed for 45 min to a 900-MHz electromagnetic field (1 and 3.5 W/kg) were compared to those of sham-exposed and cage-control rats. There were no differences among exposed, sham, and cage-control rats in the two spatial learning tasks. The discussion focuses on the potential reasons that led previous studies to conclude that learning deficits do occur after electromagnetic field exposure.

Edelstyn N, Oldershaw A. The acute effects of exposure to the electromagnetic field emitted by mobile phones on human attention. *Neuroreport* 13(1):119-121, 2002. The aim of our study was to investigate the effects of acute mobile phone exposure on a range of tasks which tapped capacity and processing speed within the attentional system. Thirty-eight healthy

volunteers were randomly assigned to either an experimental group which was exposed to a connected mobile phone or a control group in which the mobile phone was switched off. Subjects remained blind to mobile phone status throughout duration of study. The experimental group were exposed to an electromagnetic field emitted by a 900 MHz mobile phone for 30 min. Cognitive performance was assessed at three points (prior to mobile phone exposure, at 15 and 30 min post-exposure) using six cognitive neuropsychological tests (digit span and spatial span forwards and backwards, serial subtraction and verbal fluency). Significant differences between the two groups were evident after 5 min on two tests of attentional capacity (digit span forwards and spatial span backwards) and one of processing speed (serial subtraction). In all three instances, performance was facilitated following mobile phone exposure. No deficits were evident. These findings are discussed in terms of possible functional and neuroanatomical bases.

Eulitz, C, Ullsperger, P, Freude, G, Elbert, T, Mobile phones modulate response patterns of human brain activity. *Neuroreport* 9(14):3229-3232, 1998.

Mobile phones emit a pulsed high-frequency electromagnetic field (PEMF) which may penetrate the scalp and the skull. Increasingly, there is an interest in the interaction of this pulsed microwave radiation with the human brain. Our investigations show that these electromagnetic fields alter distinct aspects of the brain's electrical response to acoustic stimuli. More precisely, our results demonstrate that aspects of the induced but not the evoked brain activity during PEMF exposure can be different from those not influenced by PEMF radiation. This effect appears in higher frequency bands when subjects process task-relevant target stimuli but was not present for irrelevant standard stimuli. As the induced brain activity in higher frequency bands has been proposed to be a correlate of coherent high-frequency neuronal activity, PEMF exposure may provide means to systematically alter the pattern fluctuations in neural mass activity.

Finnie JW, Blumbergs PC, Manavis J, Utteridge TD, Gebiski V, Swift JG, Vernon-Roberts B, Kuchel TR. Effect of global system for mobile communication (gsm)-like radiofrequency fields on vascular permeability in mouse brain. *Pathology* 33(3):338-340, 2001. The effect of global system for mobile communication (GSM) radiofrequency fields on vascular permeability in the brain was studied using a purpose-designed exposure system at 898.4 MHz. Mice (n= 30) were given a single far field, whole body exposure for 60 minutes at a specific absorption rate of 4 W/kg. Control mice were also sham-exposed (n = 10) or permitted free movement in a cage (n = 10) to exclude any stress-related effects. Vascular permeability changes were detected using albumin immunohistochemistry and the efficacy of this vascular tracer was confirmed with a positive control group exposed to a clostridial toxin known to increase vascular permeability in the brain. No significant difference in albumin extravasation was

detected between any of the groups at the light microscope level using the albumin marker.

Finnie JW, Blumbergs PC, Manavis J, Utteridge TD, Gebiski V, Davies RA, Vernon-Roberts B, Kuchel TR. Effect of long-term mobile communication microwave exposure on vascular permeability in mouse brain. *Pathology* 34(4):344-347, 2002. AIMS: To study the effect of long-term exposure to global system for mobile communication (GSM) radiofrequency fields on vascular permeability in murine brains. METHODS: Using a purpose-designed exposure system at 900 MHz, mice were given a 60-minute far-field, whole body exposure on each of 5 days per week for 104 weeks at specific absorption rates (SAR) of 0.25, 1.0, 2.0 and 4.0 W/kg. Control mice were sham-exposed or permitted free movement in a cage to evaluate any stress-related effects. Albumin immunohistochemistry was used to detect increased vascular permeability and the efficacy of the vascular tracer was confirmed with a positive control group exposed to a clostridial toxin known to increase vascular permeability in the brain. RESULTS: In all exposed and control groups, albumin extravasation was minimal, often leptomeningeal, and was deemed insignificant as a maximum of three capillaries or venules in a given brain showed leakage from the very many blood vessels present in the three coronal brain sections. CONCLUSIONS: These results suggest that prolonged exposure to mobile telephone-type radiation produces negligible disruption to blood-brain barrier integrity at the light microscope level using endogenous albumin as a vascular tracer.

French PW, Donnellan M, McKenzie DR, Electromagnetic radiation at 835 MHz changes the morphology and inhibits proliferation of a human astrocytoma cell line. *Bioelectrochem Bioenerg* 43:13-18, 1997.

A human astrocytoma cell line, U-87 MG, was exposed to 835 MHz electromagnetic radiation for 20 min, 3 times per day for 7 days, at a power density of either $40 \pm 15 \text{ mWcm}^{-2}$ or $8.1 \pm 3 \text{ mWcm}^{-2}$. At the low power density, it was observed that the rate of DNA synthesis decreased, and that the cells flattened and spread out in comparison to unexposed culture. At 40 mWcm^{-2} , there were no effects seen on cell proliferation, but alteration in cell morphology included increased cell spreading and also the appearance of actin-containing blebs at localized sites on the membrane. It is hypothesised that 835 MHz radiation at low power density may be affecting a signal transduction pathway involved in cell proliferation.

Freude, G, Ullsperger, P, Eggert, S, Ruppe, I, Effects of microwaves emitted by cellular phones on human slow brain potentials. *Bioelectromagnetics* 19(6):384-387, 1998.

The influence of electromagnetic fields (EMF) emitted by cellular phones on preparatory slow brain potentials (SP) was studied in two different experimental

tasks: In the first, healthy male human subjects had to perform simple self-paced finger movements to elicit a Bereitschaftspotential; in the second, they performed a complex and cognitive demanding visual monitoring task (VMT). Both tasks were performed with and without EMF exposure in counterbalanced order. Whereas subjects' performance did not differ between the EMF exposure conditions, SP parameters were influenced by EMF in the VMT: EMF exposure effected a significant decrease of SPs at central and temporo-parieto-occipital brain regions, but not at the frontal one. In the simple finger movement task, EMF did not affect the Bereitschaftspotential.

Freude, G, Ullsperger, P, Eggert, S, Ruppe, I, Microwaves emitted by cellular telephones affect human slow brain potentials. *Eur J Appl Physiol* 81(1-2):18-27, 2000.

The influence of electromagnetic fields (EMF) emitted by cellular telephones on preparatory slow brain potentials (SP) was studied in two experiments, about 6 months apart. In the first experiment, a significant decrease of SP was found during exposure to EMF in a complex visual monitoring task (VMT). This effect was replicated in the second experiment. In addition to the VMT, EMF effects on SP were analysed in two further, less demanding tasks: in a simple finger movement task to elicit a Bereitschaftspotential (BP) and in a two-stimulus task to elicit a contingent negative variation (CNV). In comparison to the VMT, no significant main EMF effects were found in BP and CNV tasks. The results accounted for a selective EMF effect on particular aspects of human information processing, but did not indicate any influence on human performance, well-being and health.

Fritze K, Wiessner C, Kuster N, Sommer C, Gass P, Hermann DM, Kiessling M, Hossmann KA, Effect of global system for mobile communication microwave exposure on the genomic response of the rat brain. *Neuroscience* 81(3):627-639, 1997.

The acute effect of global system for mobile communication (GSM) microwave exposure on the genomic response of the central nervous system was studied in rats by measuring changes in the messenger RNAs of hsp70, the transcription factor genes c-fos and c-jun and the glial structural gene GFAP using in situ hybridization histochemistry. Protein products of transcription factors, stress proteins and marker proteins of astroglial and microglial activation were assessed by immunocytochemistry. Cell proliferation was evaluated by bromodeoxyuridine incorporation. A special GSM radiofrequency test set, connected to a commercial cellular phone operating in the discontinuous transmission mode, was used to simulate GSM exposure. The study was conducted at time averaged and brain averaged specific absorption rates of 0.3 W/kg (GSM exposure), 1.5 W/kg (GSM exposure) and 7.5 W/kg (continuous wave exposure), respectively. Immediately after exposure, in situ hybridization

revealed slight induction of hsp70 messenger RNA in the cerebellum and hippocampus after 7.5 W/kg exposure, but not at lower intensities. A slightly increased expression of c-fos messenger RNA was observed in the cerebellum, neocortex and piriform cortex of all groups subjected to immobilization, but no differences were found amongst different exposure conditions. C-jun and GFAP messenger RNAs did not increase in any of the experimental groups. 24 h after exposure, immunocytochemical analysis of FOS and JUN proteins (c-FOS, FOS B, c-JUN JUN B, JUN D), of HSP70 or of KROX-20 and -24 did not reveal any alterations. Seven days after exposure, neither increased cell proliferation nor altered expression of astroglial and microglial marker proteins were observed. In conclusion, acute high intensity microwave exposure of immobilized rats may induce some minor stress response but does not result in lasting adaptive or reactive changes of the brain.

Fritze K, Sommer C, Schmitz B, Mies G, Hossmann KA, Kiessling M, Wiessner C, Effect of global system for mobile communication (GSM) microwave exposure on blood-brain barrier permeability in rat. *Acta Neuropathol (Berl)* 94(5):465-470, 1997.

We investigated the effects of global system for mobile communication (GSM) microwave exposure on the permeability of the blood-brain barrier using a calibrated microwave exposure system in the 900 MHz band. Rats were restrained in a carousel of circularly arranged plastic tubes and sham-exposed or microwave irradiated for a duration of 4 h at specific brain absorption rates (SAR) ranging from 0.3 to 7.5 W/kg. The extravasation of proteins was assessed either at the end of exposure or 7 days later in three to five coronal brain slices by immunohistochemical staining of serum albumin. As a positive control two rats were subjected to cold injury. In the brains of freely moving control rats (n = 20) only one spot of extravasated serum albumin could be detected in one animal. In the sham-exposed control group (n = 20) three animals exhibited a total of 4 extravasations. In animals irradiated for 4 h at SAR of 0.3, 1.5 and 7.5 W/kg (n = 20 in each group) five out of the ten animals of each group killed at the end of the exposure showed 7, 6 and 14 extravasations, respectively. In the ten animals of each group killed 7 days after exposure, the total number of extravasations was 2, 0 and 1, respectively. The increase in serum albumin extravasations after microwave exposure reached significance only in the group exposed to the highest SAR of 7.5 W/kg but not at the lower intensities. Histological injury was not observed in any of the examined brains. Compared to other pathological conditions with increased blood-brain barrier permeability such as cold injury, the here observed serum albumin extravasations are very modest and, moreover, reversible. Microwave exposure in the frequency and intensity range of mobile telephony is unlikely to produce pathologically significant changes of the blood-brain barrier permeability.

Gos P, Eicher B, Kohli J, Heyer WD, No mutagenic or recombinogenic effects of mobile phone fields at 900 MHz detected in the yeast

saccharomyces cerevisiae. *Bioelectromagnetics* 21(7):515-523, 2000. Both actively growing and resting cells of the yeast *Saccharomyces cerevisiae* were exposed to 900-MHz fields that closely matched the Global System for Mobile Communication (GSM) pulsed modulation format signals for mobile phones at specific absorption rates (SAR) of 0.13 and 1.3 W/kg. Two identical anechoic test chambers were constructed to perform concurrent control and test experiments under well-controlled exposure conditions. Using specific test strains, we examined the genotoxic potential of mobile phone fields, alone and in combination, with a known genotoxic compound, the alkylating agent methyl methanesulfonate. Mutation rates were monitored by two test systems, a widely used gene-specific forward mutation assay at CAN1 and a wide-range assay measuring the induction of respiration-deficient (petite) clones that have lost their mitochondrial function. In addition, two further assays measured the recombinogenic effect of mobile phone fields to detect possible effects on genomic stability: First, an intrachromosomal, deletion-formation assay previously developed for genotoxic screening; and second, an intragenic recombination assay in the ADE2 gene. Fluctuation tests failed to detect any significant effect of mobile phone fields on forward mutation rates at CAN1, on the frequency of petite formation, on rates of intrachromosomal deletion formation, or on rates of intragenic recombination in the absence or presence of the genotoxic agent methyl methanesulfonate.

Goswami PC, Albee LD, Parsian AJ, Baty JD, Moros EG, Pickard WF, Roti Roti JL, Hunt CR, Proto-oncogene mRNA levels and activities of multiple transcription factors in C3H 10T 1/2 murine embryonic fibroblasts exposed to 835.62 and 847.74 MHz cellular phone communication frequency radiation. *Radiat Res* 151(3):300-309, 1999.

This study was designed to determine whether two differently modulated radiofrequencies of the type generally used in cellular phone communications could elicit a general stress response in a biological system. The two modulations and frequencies studied were a frequency-modulated continuous wave (FMCW) with a carrier frequency of 835.62 MHz and a code division multiple-access (CDMA) modulation centered on 847.74 MHz. Changes in proto-oncogene expression, determined by measuring Fos, Jun, and Myc mRNA levels as well as by the DNA-binding activity of the AP1, AP2 and NF-kappaB transcription factors, were used as indicators of a general stress response. The effect of radiofrequency exposure on proto-oncogene expression was assessed (1) in exponentially growing C3H 10T 1/2 mouse embryo fibroblasts during their transition to plateau phase and (2) during transition of serum-deprived cells to the proliferation cycle after serum stimulation. Exposure of serum-deprived cells to 835.62 MHz FMCW or 847.74 MHz CDMA microwaves (at an average specific absorption rate, SAR, of 0.6 W/kg) did not significantly change the kinetics of proto-oncogene expression after serum stimulation. Similarly, these exposures did not affect either the Jun and Myc mRNA levels or the DNA-binding activity of AP1, AP2 and NF-kappaB in exponential cells during transit to plateau-phase

growth. Therefore, these results suggest that the radiofrequency exposure is unlikely to elicit a general stress response in cells of this cell line under these conditions. However, statistically significant increases (approximately 2-fold, $P = 0.001$) in Fos mRNA levels were detected in exponential cells in transit to the plateau phase and in plateau-phase cells exposed to 835.62 MHz FMCW microwaves. For 847.74 MHz CDMA exposure, the increase was 1.4-fold ($P = 0.04$). This increase in Fos expression suggests that expression of specific genes could be affected by radiofrequency exposure.

Haarala C, Bjornberg L, Ek M, Laine M, Revonsuo A, Koivisto M, Hamalainen H. Effect of a 902 MHz electromagnetic field emitted by mobile phones on human cognitive function: A replication study.

***Bioelectromagnetics* 24(4):283-288, 2003.** Our study was a replication and extension with methodological improvements to a previous study on effects of the electromagnetic field (EMF) emitted by a 902 MHz mobile phone on human cognitive functioning. Improvements on the previous study included multicentre testing and a double blind design. A total of 64 subjects (32 men and 32 women) in two independent laboratories performed a battery of 9 cognitive tasks twice: while the EMF was on and while it was off. Reaction times (RTs) and accuracy were recorded. The order of exposure and tasks was counterbalanced across subjects and gender. There were no statistically significant differences in performance between genders or laboratories. Although the RTs and the accuracy of answers were very similar to those of our previous study, our previous results were not replicated. We concluded that EMF had no effect on RTs or on the accuracy of the subjects' answers. Further, our results indicate that our EMF had no immediate effect on human cognitive functioning or that such effects are so small that they are observed on behavior only occasionally.

Hardell L, Hansson Mild K, Pahlson A, Hallquist A, Ionizing radiation, cellular telephones and the risk of brain tumours. *Europ J Cancer Prevent* 10:523-529, 2001.

A case-control study on brain tumours included 233 patients aged 20-80 years and alive at the study time. They had histopathologically verified brain tumour and lived in the Upsala-Orebro region (1994-1996) or in Stockholm region (1995-1996). Two matched controls to each case were selected from the Swedish Population Register. Two hundred and nine cases (90%) and 425 controls (91%) answered the questionnaire. Results are presented for the whole study group, as given here, and for malignant and benign tumours separately. For workers in the chemical industry the odds ratio (OR) was 4.10, 95% confidence interval (95% CI) 1.25-13.4 and laboratory workers OR 3.21, 95%CI 1.16-8.85. Radiotherapy of the head and neck region gave OR 3.61, 95% CI 0.65-19.9. Medical diagnostic X-ray of the same area yielded OR 1.64, 95% CI 1.04-2.58. Work as a physician gave OR 6.00, 95% CI 0.62-57.7. All three cases had worked with fluoroscopy. Ipsilateral (same side) use of a cellular telephone increased the risk of tumours in

the temporal, temporo-parietal and occipital areas, with OR 2.42, 95% CI 0.97-6.05 (i.e., the anatomical areas with highest exposure to microwaves from a mobile phone).

Hardell, L, Nasman, A, Pahlson, A, Hallquist, A, Hansson Mild, K, Use of cellular telephones and the risk for brain tumours: A case-control study. *Int J Oncol* 15(1):113-116, 1999.

The use of cellular telephones has increased dramatically during the 1990's in the world. In the 1980's the analogue NMT system was used whereas the digital GSM system was introduced in early 1990's and is now the preferred system. Case reports of brain tumours in users initiated this case-control study on brain tumours and use of cellular telephones. Also other exposures were assessed. All cases, both males and females, with histopathologically verified brain tumour living in Uppsala-Orebro region (1994-96) and Stockholm region (1995-96) aged 20-80 at the time of diagnosis and alive at start of the study were included, 233 in total. Two controls to each case were selected from the Swedish Population Register matched for sex, age and study region. Exposure was assessed by questionnaires supplemented over the phone. The analyses were based on answers from 209 (90%) cases and 425 (91%) controls. Use of cellular telephone gave odds ratio (OR) = 0.98 with 95% confidence interval (CI) = 0.69-1.41. For the digital GSM system OR = 0.97, CI = 0.61-1.56 and for the analogue NMT system OR = 0.94, CI = 0.62-1.44 were calculated. Dose-response analysis and using different tumour induction periods gave similar results. Non-significantly increased risk was found for tumour in the temporal or occipital lobe on the same side as a cellular phone had been used, right side OR = 2.45, CI = 0.78-7.76, left side OR = 2.40, CI = 0.52-10.9 Increased risk was found only for use of the NMT system. For GSM use the observation time is still too short for definite conclusions. An increased risk for brain tumour in the anatomical area close to the use of a cellular telephone should be especially studied in the future.

Hardell L, Nasman A, Pahlson A, Hallquist A, Case-Control Study on Radiology Work, Medical X-ray Investigations, and Use of Cellular Telephones as Risk Factors for Brain Tumors. *Medscape General Medicine* May 4, 2000.

Abstract

Context. Ionizing radiation is a well-established risk factor for brain tumors. During recent years, microwave exposure from the use of cellular telephones has been discussed as a potential risk factor. **Objective.** To determine risk factors for brain tumors. **Design.** A case-control study, with exposure assessed by questionnaires. **Participants.** A total of 233 currently living men and women, aged 20 to 80 years, were included. The case patients had histopathologically verified brain tumors and lived in the Uppsala-Orebro region (1994-1996) or the Stockholm region (1995-1996). Two matched controls to each case were selected from the Swedish Population Register. **Main Outcome Measures.**

Ionizing radiation and use of cellular telephones as risk factors for brain tumors. **Results.** A total of 209 cases (90%) and 425 controls (91%) answered the questionnaire. Work as a physician yielded an odds ratio (OR) of 6.00, with a 95% confidence interval (CI) of 0.62 to 57.7. All three case patients had worked with fluoroscopy. Radiotherapy of the head and neck region yielded an OR of 3.61 (95% CI, 0.65-19.9). Medical diagnostic x-ray examination of the same area yielded an OR of 2.10 (95% CI, 1.25-3.53), with a tumor induction period of 5 years or more. Chemical industry work yielded an OR of 4.10 (95% CI, 1.25-13.4), and laboratory work yielded an OR of 3.21 (95% CI, 1.16-8.85). Ipsilateral use of cellular telephones increased the risk for tumors in the temporal, temporoparietal, and occipital lobes (OR, 2.42; 95% CI, 0.97-6.05), ie, the anatomic areas with highest exposure to microwaves from a mobile telephone. The result was further strengthened (OR, 2.62; 95% CI, 1.02-6.71) in a multivariate analysis that included laboratory work and medical diagnostic x-ray investigations of the head and neck. **Conclusion.** Exposure to ionizing radiation, work in laboratories, and work in the chemical industry increased the risk of brain tumors. Use of a cellular telephone was associated with an increased risk in the anatomic area with highest exposure.

Hardell L, Hallquist A, Hansson Mild K, Carlberg M, Pahlson A, Lilja A. cellular and cordless telephones and the risk for brain tumours. *Europ J Cancer Prevent* 11:377-386, 2002.

Microwave exposure from the use of cellular telephones has been discussed in recent years as a potential risk factor for brain tumours. We included in a case-control study 1617 patients aged 20-80 years of both sexes with brain tumour diagnosed between 1 January 1997 and 30 June 2000. They were alive at the study time and had histopathologically verified brain tumour. One matched control to each case was selected from the Swedish Population Register. The study area was the Uppsala-Orebro, Stockholm, Linköping and Göteborg medical regions of Sweden. Exposure was assessed by a questionnaire that was answered by 1429 (88%) cases and 1470 (91%) controls. In total, use of analogue cellular telephones gave an increased risk with an odds ratio (OR) of 1.3 (95% confidence interval (CI) 1.02-1.6). With a tumour induction period of >10 years the risk increased further; OR 1.8 (95% CI 1.1-2.9). No clear association was found for digital or cordless telephones. With regard to the anatomical area of the tumour and exposure to microwaves, the risk was increased for tumours located in the temporal area on the same side of the brain that was used during phone calls; for analogue cellular telephones the OR was 2.5 (95% CI 1.3-4.9). Use of a telephone on the opposite side of the brain was not associated with an increased risk for brain tumours. With regard to different tumour types, the highest risk was for acoustic neurinoma (OR 3.5, 95% CI 1.8-6.8) among analogue cellular telephone users.

Hardell L, Mild KH, Carlsberg M. Case-control study on the use of cellular and cordless phones and the risk for malignant brain tumours. *Int. J. Radiat. Biol.* 78:931-936, 2002.

Purpose: To investigate the use of cellular and cordless phones and the risk for malignant brain tumours.

Materials and Methods: A case-control study was performed on 649 patients aged 20-80 years of both sexes with malignant brain tumour diagnosed from 1 January 1997 to 30 June 2000. All patients were alive during the time of the study and had histopathology verified brain tumours. One matched control to each case was selected from the Swedish Population Register. The study area was the Uppsala-Orebro, Stockholm, Linköping and Göteborg medical regions of Sweden.

Results: Exposure was assessed by a questionnaire answered by 588 (91%) cases and 581 (90%) controls. Phone usage was defined as 'ever use' and usage starting 1 year before diagnosis was disregarded. Overall, no significantly increased risks were found: analogue cellular phones yielded an odds ratio (OR) = 1.13, 95% confidence interval (CI) = 0.82-1.57, digital cellular phones OR = 1.13, CI = 0.86-1.48, and cordless phones OR = 1.13, CI = 0.85-1.50. For ipsilateral (same side) radiofrequency exposure, analogue mobile phones gave OR = 1.85, CI = 1.16-2.96, for all malignant brain tumours. For astrocytoma, this risk was OR = 1.95, CI = 1.12-3.39. For all malignant brain tumours, digital mobile phones yielded OR = 1.59, CI = 1.05-2.41, and cordless phones yielded OR = 1.46, CI = 0.96-2.23, in the analysis of ipsilateral exposure.

Conclusion: The ipsilateral use of an analogue cellular phone yielded a significantly increased risk for malignant brain tumours.

Hardell L, Mild KH, Carlberg M. Further aspects on cellular and cordless telephones and brain tumours. *Int J Oncol* 22(2):399-407, 2003.

We included in a case-control study on brain tumours and mobile and cordless telephones 1,617 patients aged 20-80 years of both sexes diagnosed during January 1, 1997 to June 30, 2000. They were alive at the study time and had histopathology verified brain tumour. One matched control to each case was selected from the Swedish Population Register. The study area was the Uppsala-Orebro, Stockholm, Linköping and Göteborg medical regions of Sweden.

Exposure was assessed by a questionnaire that was answered by 1,429 (88%) cases and 1,470 (91%) controls. In total use of analogue cellular telephones gave an increased risk with odds ratio (OR)=1.3, 95% confidence interval (CI)=1.04-1.6, whereas digital and cordless phones did not overall increase the risk significantly. Ipsilateral use of analogue phones gave OR=1.7, 95% CI=1.2-2.3, digital phones OR=1.3, 95% CI=1.02-1.8 and cordless phones OR=1.2, 95% CI=0.9-1.6. The risk for ipsilateral use was significantly increased for astrocytoma for all studied phone types, analogue phones OR=1.8, 95% CI=1.1-3.2, digital phones OR=1.8, 95% CI=1.1-2.8, cordless phones OR=1.8, 95% CI=1.1-2.9.

Use of a telephone on the opposite side of the brain was not associated with a significantly increased risk for brain tumours. Regarding anatomical area of the

tumour and exposure to microwaves, the risk was increased for tumours located in the temporal area on the same side of the brain that was used during phone calls, significantly so for analogue cellular telephones OR=2.3, 95% CI=1.2-4.1. For acoustic neurinoma OR=4.4, 95% CI=2.1-9.2 was calculated among analogue cellular telephone users. When duration of use was analysed as a continuous variable in the total material, the risk increased per year for analogue phones with OR=1.04, 95% CI=1.01-1.08. For astrocytoma and ipsilateral use the trend was for analogue phones OR=1.10, 95% CI=1.02-1.19, digital phones OR=1.11, 95% CI=1.01-1.22, and cordless phones OR=1.09, 95% CI=1.01-1.19. There was a tendency of a shorter tumour induction period for ipsilateral exposure to microwaves than for contralateral, which may indicate a tumour promotor effect.

Hardell L, Mild KH, Sandstrom M, Carlberg M, Hallquist A, Pahlson A. Vestibular schwannoma, tinnitus and cellular telephones. *Neuroepidemiol* 22:124-129, 2003.

Cases with tinnitus after using analogue cellular telephones are presented. An increased odds ratio of 3.45, 95% confidence interval (CI) 1.77-6.76, was found for vestibular schwannoma (VS) associated with the use of analogue cell phones. During the time period 1960-1998, the age-standardized incidence of VS in Sweden significantly increased yearly by +2.53% (CI 1.71-3.35). A significant increase in the incidence of VS was only found for the latter of the two time period 1960-1979 and 1980-1998. For all other brain tumors taken together, the incidence significantly increased yearly by +0.80% (CI 0.59-1.02) for the time period 1960-1998, although the increase was only significant for benign tumors other than VS during 1960-1979.

Harvey C, French PW, Effects on protein kinase C and gene expression in a human mast cell line, HMC-1, following microwave exposure. *Cell Biol Int* 23(11):739-748, 2000.

We used a resonant cavity which delivered a continuous wave exposure at 864.3 MHz at an average specific absorption rate (SAR) of 7 W/kg to determine non-thermal biological effects of microwave exposure. A human mast cell line, HMC-1, was used as the biological target. Cells were given three exposures each of 20-min duration daily for 7 days. The temperature of the cell culture medium during the exposure fell to 26.5 degrees C. Effects were seen on localization of protein kinase C, and expression of three genes of 588 screened. The affected genes included the proto-oncogene c-kit, the transcription factor Nucleoside diphosphate kinase B and the apoptosis-associated gene DAD-1. Stress response genes were variably upregulated. No significant effect on morphology or on F-actin distribution was detected. We conclude that low-power microwave exposure may act on HMC-1 cells by altering gene expression via a mechanism involving activation of protein kinase C, and at temperatures well below those known to induce a heat shock response.

Heikkinen P, Kosma VM, Hongisto T, Huuskonen H, Hyysalo P, Komulainen H, Kumlin T, Lahtinen T, Lang S, Puranen L, Juutilainen J. Effects of Mobile Phone Radiation on X-Ray-Induced Tumorigenesis in Mice. *Radiat Res* 156(6):775-785, 2001. The increased use of mobile phones has raised the question of possible health effects of such devices, particularly the risk of cancer. It seems unlikely that the low-level radiofrequency (RF) radiation emitted by them would damage DNA directly, but its ability to act as a tumor promoter is less well characterized. In the current study, we evaluated the effect of low-level RF radiation on the development of cancer initiated in mice by ionizing radiation. Two hundred female CBA/S mice were randomized into four equal groups at the age of 3 to 5 weeks. The mice in all groups except the cage-control group were exposed to ionizing radiation at the beginning of the study and then to RF radiation for 1.5 h per day, 5 days a week for 78 weeks. One group was exposed to continuous NMT (Nordic Mobile Telephones)-type frequency-modulated RF radiation at a frequency of 902.5 MHz and a nominal average specific absorption rate (SAR) of 1.5 W/kg. Another group was exposed to pulsed GSM (Global System for Mobile)-type RF radiation (carrier-wave frequency 902.4 MHz, pulse frequency 217 Hz) at a nominal average SAR of 0.35 W/kg. The control animals were sham-exposed. Body weight, clinical signs, and food and water consumption were recorded regularly. Hematological examinations and histopathological analyses of all lesions and major tissues were performed on all animals. The RF-radiation exposures did not increase the incidence of any neoplastic lesion significantly. We conclude that the results do not provide evidence for cancer promotion by RF radiation emitted by mobile phones.

Heikkinen P, Kosma VM, Alhonen L, Huuskonen H, Komulainen H, Kumlin T, Laitinen JT, Lang S, Puranen L, Juutilainen J. Effects of mobile phone radiation on UV-induced skin tumourigenesis in ornithine decarboxylase transgenic and non-transgenic mice. *Int J Radiat Biol* 79(4):221-233, 2003. Purpose: The effects of low-level radiofrequency radiation (RFR) on ultraviolet (UV)-induced skin tumorigenesis were evaluated in ornithine decarboxylase (ODC) and non-transgenic mice. Materials and methods: Transgenic female mice over-expressing the human ODC gene and their non-transgenic littermates (20 animals in the cage control group, and 45-49 animals in the other groups) were exposed for 52 weeks to UV radiation or a combination of UV radiation and pulsed RFR. The UV dose was 240 Jm⁻² (1.2 x human minimum erythema dose) delivered three times a week. One group of animals was exposed to Digital Advanced Mobile Phone System (DAMPS)-type RFR, the other group to Global System for Mobile (GSM)-type RFR at a nominal average specific absorption rate of 0.5 W kg⁻¹, 1.5 h day⁻¹, for 5 days a week. The skin was carefully palpated weekly for macroscopic tumours. Histopathological analyses of all skin lesions and of a specified dorsal skin area were performed on all animals. Results: UV exposure resulted in development of macroscopic skin tumours in 11.5 and 36.8% of non-transgenic and transgenic animals, respectively. The RFR exposures did not give a statistically significant effect on the development of skin

tumours in either transgenic or non-transgenic animals, or in combined analysis, but tumour development appeared slightly accelerated especially in non-transgenic animals. No effects of RFR exposures were found on excretion of 6-hydroxymelatonin sulphate into urine or on polyamine levels in dorsal skin. Conclusion: RFR exposures did not significantly enhance skin tumourigenesis. However, the slightly accelerated tumour development may warrant further evaluation.

Hietanen M, Kovala T, Hamalainen AM, Human brain activity during exposure to radiofrequency fields emitted by cellular phones. *Scand J Work Environ Health* 26(2):87-92, 2000.

OBJECTIVES: The aim of this study was to explore the possible influence of radiofrequency (RF) radiation exposure on human brain function. **METHODS:** The electroencephalographic (EEG) activity of 19 volunteers was quantitatively analyzed. Ten of the subjects were men (28-48 years of age) and 9 were women (32-57 years of age). The sources of exposure were 5 different cellular phones (analogue and digital models) operating at a frequency of 900 MHz or 1800 MHz. The EEG activity was recorded in an awake, closed-eyes situation. Six 30-minute experiments, including 1 sham exposure, were made for each subject. The duration of a real exposure phase was 20 minutes. **RESULTS:** Exposure to one of the phones caused a statistically significant change in the absolute power at the delta band of the EEG recording. However, no difference was seen in the relative power of the same band, and no changes occurred during exposure to other phones at any frequency bands. **CONCLUSIONS:** The findings of this study suggest that exposure to radiofrequency fields emitted by cellular phones has no abnormal effects on human EEG activity. The observed difference in 1 parameter was probably caused by statistical chance.

Hietanen M, Hämäläinen A-M, Husman T. Hypersensitivity symptoms associated with exposure to cellular telephones: No causal link. *Bioelectromagnetics* 23:264-270, 2002.

The hypothesis that there exist hypersensitive persons who perceive subjective symptoms from radiofrequency (RF) fields emitted by hand held mobile phones (cellular phones) was tested using double blind provocation experiments. We also tested whether sensitive subjects are able to determine whether the phone is on or off by sensing RF fields. The study group consisted of 20 volunteers (13 women and 7 men) who reported themselves as being sensitive to cellular phones. The RF exposure sources were one analogue NMT phone (900 MHz) and two digital GSM phones (900 and 1800 MHz). The duration of a test session was 30 min, and three or four sessions were performed in random order for each subject during 1 day. The subjects were asked to report symptoms or sensations as soon as they perceived any abnormal feelings. In addition, the subjects' blood pressure, heart rate, and breathing frequency were monitored every 5 min. The results of the study indicated that various symptoms were reported, and most of

them appeared in the head region. However, the number of reported symptoms was higher during sham exposure than during real exposure conditions. In addition, none of the test persons could distinguish real RF exposure from sham exposure. Hence, we conclude that adverse subjective symptoms or sensations, though unquestionably perceived by the test subjects, were not produced by cellular phones.

Higashikubo R, Culbreth VO, Spitz DR, LaRegina MC, Pickard WF, Straube WL, Moros EG, Roti JL, Radiofrequency electromagnetic fields have no effect on the in vivo proliferation of the 9L brain tumor. *Radiat Res* 152(6):665-671, 1999.

The intracranial 9L tumor model was used to determine if exposure to a radiofrequency (RF) electromagnetic field similar to those used in cellular telephone has any effects on the growth of a central nervous system tumor. Fischer 344 rats implanted with different numbers of 9L gliosarcoma cells were exposed to 835.62 MHz frequency-modulated continuous wave (FMCW) or 847.74 MHz code division multiple access (CDMA) RF field with nominal slot-average specific absorption rates in the brain of 0.75 +/- 0.25 W/kg. The animals were exposed to the RF field for 4 h a day, 5 days a week starting 4 weeks prior to and up to 150 days after the implantation of tumor cells. Among sham-exposed animals injected with 2 to 10 viable cells (group 1), the median survival was 70 days, with 27% of the animals surviving at 150 days. The median survival length and final survival fraction for animals injected with 11 to 36 viable cells (group 2) were 52 days and 14%, respectively, while the values for those injected with 37 to 100 cells (group 3) were 45 days and 0%. The animals exposed to CDMA or FMCW had similar survival parameters, and the statistical comparison of the survival curves for each of the groups 1, 2 and 3 showed no significant differences compared to sham-exposed controls.

Higashikubo R, Ragouzis M, Moros EG, Straube WL, Roti Roti JL. Radiofrequency Electromagnetic Fields do not Alter the Cell Cycle Progression of C3H 10T and U87MG Cells. *Radiat Res* 156(6):786-795, 2001.

The effects of exposure to radiofrequency electromagnetic fields (RF EMFs) on cell cycle progression of mouse fibroblasts C3H 10T(1/2) and human glioma U87MG cells were determined by the flow cytometric bromodeoxyuridine pulse-chase method. Cells were exposed to a frequency-modulated continuous wave at 835.62 MHz or a code division multiple access RF EMF centered on 847.74 MHz at an average specific absorption rate of 0.6 W/kg. Five cell cycle parameters, including the transit of cells through G(1), G(2) and S phase and the probability of cell division, were examined immediately after the cells were placed in the fields or after they had been kept in the fields for up to 100 h. The only significant change observed in the study was that associated with C3H 10T(1/2) cell cultures moving into plateau phase toward the later times in the long-exposure experiment. No changes in the cell cycle parameters were observed in cells exposed to either mode of RF EMFs when compared to sham-exposed cells

in either of the cell lines studied during the entire experimental period. The results show that exposure to RF EMFs, at the frequencies and power tested, does not have any effect on cell progression in vitro.

Hladky, A, Musil, J, Roth, Z, Urban, P, Blazkova, V, Acute effects of using a mobile phone on CNS functions. *Cent Eur J Public Health* 7(4):165-167. 1999.

Twenty volunteers participated in two experiments exploring the acute effects of using the mobile phone Motorola GSM 8700 on the functions of the CNS. When speaking (5 minutes reading a text from daily newspapers) the electromagnetic fields from the mobile apparatus did not affect the visual evoked potentials. Also a 6-min exposure did not reveal any effect of electromagnetic fields on the results in two tests (memory and attention) performed while speaking into the mobile. On the other hand the phone call itself strongly influenced the performance in a secondary task applying a test of switching attention which is a good model for driving a car. The response and decision speed were significantly worse. This is a proof that even a slight psychological stress involved in calling while driving can be a great risk.

Hocking, B, Preliminary report: symptoms associated with mobile phone use. *Occup Med (Lond)*;48(6):357-360, 1998.

Mobile phone use is ubiquitous, although the alleged health effects of low level radio-frequency radiation (RFR) used in transmission are contentious. Following isolated reports of headache-like symptoms arising in some users, a survey has been conducted to characterize the symptoms sometimes associated with mobile phone usage. A notice of interest in cases was placed in a major medical journal and this was publicized by the media. Respondents were interviewed by telephone using a structured questionnaire. Forty respondents from diverse occupations described unpleasant sensations such as a burning feeling or a dull ache mainly occurring in the temporal, occipital or auricular areas. The symptoms often began minutes after beginning a call, but could come on later during the day. The symptoms usually ceased within an hour after the call, but could last until evening. Symptoms did not occur when using an ordinary handset, and were different from ordinary headaches. There were several reports suggestive of intra-cranial effects. Three respondents reported local symptoms associated with wearing their mobile phone on their belts. There was one cluster of cases in a workplace. Seventy-five per cent of cases were associated with digital mobile phones. Most of the respondents obtained relief by altering their patterns of telephone usage or type of phone. Cranial and other diverse symptoms may arise associated with mobile phone usage. Physicians and users alike should be alert to this. Further work is needed to determine the range of effects, their mechanism and the possible implications for safety limits of RFR.

Huber R, Graf T, Cote KA, Wittmann L, Gallmann E, Matter D, Schuderer J, Kuster N, Borbely AA, Achermann P, Exposure to pulsed high-frequency electromagnetic field during waking affects human sleep EEG. *Neuroreport* 11(15):3321-3325, 2000. The aim of the study was to investigate whether the electromagnetic field (EMF) emitted by digital radiotelephone handsets affects brain physiology. Healthy, young male subjects were exposed for 30 min to EMF (900 MHz; spatial peak specific absorption rate 1 W/kg) during the waking period preceding sleep. Compared with the control condition with sham exposure,

spectral power of the EEG in non-rapid eye movement sleep was increased. The maximum rise occurred in the 9.75-11.25 Hz and 12.5-13.25 Hz band during the initial part of sleep. These changes correspond to those obtained in a previous study where EMF was intermittently applied during sleep. Unilateral exposure induced no hemispheric asymmetry of EEG power. The present results demonstrate that exposure during waking modifies the EEG during subsequent sleep. Thus the changes of brain function induced by pulsed high-frequency EMF outlast the exposure period.

Huber R, Treyer V, Borbély AA, Schuderer J, Gottselig JM, Landolt H-P, Werth E, Berthold T, Kuster N, Buck A, Achermann P, Electromagnetic fields, such as those from mobile phones, alter regional cerebral blood flow and sleep and waking EEG. *J Sleep Res* 11: 289-295, 2002.

Usage of mobile phones is rapidly increasing, but there is limited data on the possible effects of electromagnetic field (EMF) exposure on brain physiology. We investigated the effect of EMF vs. sham control exposure on waking regional cerebral blood flow (rCBF) and on waking and sleep electroencephalogram (EEG) in humans. In Experiment 1, positron emission tomography (PET) scans were taken after unilateral head exposure to 30-min pulse-modulated 900 MHz electromagnetic field (pm-EMF). In Experiment 2, night-time sleep was polysomnographically recorded after EMF exposure. Pulse-modulated EMF exposure increased relative rCBF in the dorsolateral prefrontal cortex ipsilateral to exposure. Also, pm-EMF exposure enhanced EEG power in the alpha frequency range prior to sleep onset and in the spindle frequency range during stage 2 sleep. Exposure to EMF without pulse modulation did not enhance power in the waking or sleep EEG. We previously observed EMF effects on the sleep EEG (A. A. Borbély, R. Huber, T. Graf, B. Fuchs, E. Gallmann and P. Achermann. *Neurosci. Lett.*, 1999, 275: 207-210; R. Huber, T. Graf, K. A. Cote, L. Wittmann, E. Gallmann, D. Matter, J. Schuderer, N. Kuster, A. A. Borbély, and P. Achermann. *Neuroreport*, 2000, 11: 3321-3325), but the basis for these effects was unknown. The present results show for the first time that (1) pm-EMF alters waking rCBF and (2) pulse modulation of EMF is necessary to induce waking and sleep EEG changes. Pulse-modulated EMF exposure may provide a new, non-invasive method for modifying brain function for experimental, diagnostic and therapeutic purposes.

Achermann. *Neurosci. Lett.*, 1999, 275: 207-210; R. Huber, T. Graf, K. A. Cote, L. Wittmann, E. Gallmann, D. Matter, J. Schuderer, N. Kuster, A. A. Borbély, and P. Achermann. *Neuroreport*, 2000, 11: 3321-3325), but the basis for these effects was unknown. The present results show for the first time that (1) pm-EMF alters waking rCBF and (2) pulse modulation of EMF is necessary to induce waking and sleep EEG changes. Pulse-modulated EMF exposure may provide a new, non-invasive method for modifying brain function for experimental, diagnostic and therapeutic purposes.

Huber R, Schuderer J, Graf T, Jutz K, Borbely AA, Kuster N, Achermann P. Radio frequency electromagnetic field exposure in humans: Estimation of SAR distribution in the brain, effects on sleep and heart rate.

***Bioelectromagnetics* 24(4):262-276, 2003.** In two previous studies we demonstrated that radiofrequency electromagnetic fields (RF EMF) similar to those emitted by digital radiotelephone handsets affect brain physiology of healthy young subjects exposed to RF EMF (900 MHz; spatial peak specific absorption rate [SAR] 1 W/kg) either during sleep or during the waking period preceding sleep. In the first experiment, subjects were exposed intermittently during an 8 h nighttime sleep episode and in the second experiment, unilaterally for 30 min prior to a 3 h daytime sleep episode. Here we report an extended

analysis of the two studies as well as the detailed dosimetry of the brain areas, including the assessment of the exposure variability and uncertainties. The latter enabled a more in depth analysis and discussion of the findings. Compared to the control condition with sham exposure, spectral power of the non-rapid eye movement sleep electroencephalogram (EEG) was initially increased in the 9-14 Hz range in both experiments. No topographical differences with respect to the effect of RF EMF exposure were observed in the two experiments. Even unilateral exposure during waking induced a similar effect in both hemispheres. Exposure during sleep reduced waking after sleep onset and affected heart rate variability. Exposure prior to sleep reduced heart rate during waking and stage 1 sleep. The lack of asymmetries in the effects on sleep EEG, independent of bi- or unilateral exposure of the cortex, may indicate involvement of subcortical bilateral projections to the cortex in the generation of brain function changes, especially since the exposure of the thalamus was similar in both experiments (approx. 0.1 W/kg).

Imaida, K, Taki, M, Watanabe, S, Kamimura, Y, Ito, T, Yamaguchi, T, Ito, N, Shirai, T, The 1.5 GHz electromagnetic near-field used for cellular phones does not promote rat liver carcinogenesis in a medium-term liver bioassay. *Jpn J Cancer Res* 89(10):995-1002, 1998.

We have recently established that local exposure to a 929.2 MHz electromagnetic near-field, used for cellular phones, does not promote rat liver carcinogenesis in a medium-term bioassay system. In the present study, a 1.439 GHz electromagnetic near-field (EMF), another microwave band employed for cellular phones in Japan, was similarly investigated. Time division multiple access (TDMA) signals for the Personal Digital Cellular (PDC) Japanese cellular telephone standard system were directed to rats through a quarter-wavelength monopole antenna. Numerical dosimetry showed that the peak SARs within the liver were 1.91-0.937 W/kg, while the whole-body average specific absorption rates (SARs) were 0.680-0.453 W/kg, when the time-averaged antenna radiation power was 0.33 W. Exposure was for 90 min a day, 5 days a week, over 6 weeks, to male F344 rats given a single dose of diethylnitrosamine (200 mg/kg, i.p.) 2 weeks previously. At week 3, all rats were subjected to a two-thirds partial hepatectomy. At week 8, the experiment was terminated and the animals were killed. Carcinogenic potential was scored by comparing the numbers and areas of the induced glutathione S-transferase placental form (GST-P)-positive foci in the livers of exposed (48) and sham-exposed rats (48). Despite increased serum levels of corticosterone, adrenocorticotrophic hormone (ACTH) and melatonin, the numbers and the areas of GST-P-positive foci were not significantly altered by the exposure. These findings clearly indicated that local body exposure to a 1.439 GHz EMF, as in the case of a 929.2 MHz field, has no promoting effect on rat liver carcinogenesis in the present model.

Imaida K, Taki M, Yamaguchi T, Ito T, Watanabe S, Wake K, Aimoto A, Kamimura Y, Ito N, Shirai T, Lack of promoting effects of the electromagnetic near-field used for cellular phones (929.2 MHz) on rat liver carcinogenesis in a medium-term liver bioassay. *Carcinogenesis* 19(2):311-314, 1998. The possible cancer promotion potential of local exposure to a pulse modulated 929.2 MHz electromagnetic near-field on chemically-initiated rat liver carcinogenesis was investigated employing a medium-term bioassay. A 929.2-MHz electromagnetic near-field of time division multiple access (TDMA) signal for PDC (Personal Digital Cellular, Japanese cellular telephone standard) system was directed to rats through a quarter-wavelength monopole antenna. Maximum local specific absorption rates (SARs) on temporal average were 7.2-6.6 W/kg within the whole body and 2.0-1.7 W/kg within the liver, which was the target organ. The whole-body average SARs on temporal average were 0.80-0.58 W/kg. Temporal peak SARs had three times these values due to the duty ratio of the PDC signal. Exposure was for 90 min a day, 5 days a week, over 6 weeks. The exposure apparatus was specially designed for this experiment, to allow exposure of the lateral mid-section of the rat body to the electromagnetic near-field. Male F344 rats, 6 week-old, were initially (at week 0) given a single dose of diethylnitrosamine (DEN, 200 mg/kg body wt, i.p.). At 2 weeks later, exposure (48 rats) or sham-exposure (48 rats) was started. The exposure of electromagnetic near-fields was performed using the exposure apparatus mentioned above. At week 3, all rats were subjected to a 2/3 partial hepatectomy. At week 8 (i.e. after 6 weeks exposure or sham-exposure), the experiment was terminated and all rats were killed. Carcinogenic potential was scored by comparing the numbers and areas of the induced glutathione S-transferase placental form (GST-P) positive foci in the livers of the exposed and sham-exposed rats. A further group of 24 animals, given only DEN and partial hepatectomy, served as the controls. The numbers (no./cm²) of GST-P positive foci were 4.61 +/- 1.77, 5.21 +/- 1.92 ($P < 0.05$, versus control) and 4.09 +/- 1.47 and the areas (mm²/cm²) were 0.30 +/- 0.16, 0.36 +/- 0.21 and 0.28 +/- 0.15, for the exposed, sham-exposed and control groups, respectively. There were no significant differences between the exposed and sham-exposed groups. These findings clearly indicated that local body exposure to a 929.2-MHz field, modulated in a PDC waveform, has no significant effect on rat liver carcinogenesis under the experimental conditions employed.

Inskip PD, Tarone RE, Hatch EE, Wilcosky TC, Shapiro WR, Selker RG, Fine HA, Black PM, Loeffler JS, Linet MS, Cellular-Telephone Use and Brain Tumors. *N Engl J Med* 344(2):79-86, 2001. Background: Concern has arisen that the use of hand-held cellular telephones might cause brain tumors. If such a risk does exist, the matter would be of considerable public health importance, given the rapid increase worldwide in the use of these devices. Methods: We examined the use of cellular telephones in a case-control study of intracranial tumors of the nervous system conducted between 1994 and 1998. We enrolled 782 patients through hospitals in Phoenix, Arizona; Boston; and Pittsburgh; 489 had histologically confirmed glioma, 197 had meningioma, and 96 had acoustic neuroma. The 799 controls were patients admitted to the same hospitals as the

patients with brain tumors for a variety of nonmalignant conditions. Results: As compared with never, or very rarely, having used a cellular telephone, the relative risks associated with a cumulative use of a cellular telephone for more than 100 hours were 0.9 for glioma (95 percent confidence interval, 0.5 to 1.6), 0.7 for meningioma (95 percent confidence interval, 0.3 to 1.7), 1.4 for acoustic neuroma (95 percent confidence interval, 0.6 to 3.5), and 1.0 for all types of tumors combined (95 percent confidence interval, 0.6 to 1.5). There was no evidence that the risks were higher among persons who used cellular telephones for 60 or more minutes per day or regularly for five or more years. Tumors did not occur disproportionately often on the side of head on which the telephone was typically used. Conclusions: These data do not support the hypothesis that the recent use of hand-held cellular telephones causes brain tumors, but they are not sufficient to evaluate the risks among long-term, heavy users and for potentially long induction periods.

Ivaschuk OI, Jones RA, Ishida-Jones T, Haggren W, Adey WR, Phillips JL, Exposure of nerve growth factor-treated PC12 rat pheochromocytoma cells to a modulated radiofrequency field at 836.55 MHz: effects on c-jun and c-fos expression. *Bioelectromagnetics* 18(3):223-229, 1997.

Rat PC12 pheochromocytoma cells have been treated with nerve growth factor And then exposed to athermal levels of a packet-modulated radiofrequency field At 836.55 MHz. This signal was produced by a prototype time-domain multiple-access (TDMA) transmitter that conforms to the North American digital cellular telephone standard. Three slot average power densities were used: 0.09, 0.9, and 9 mW/cm². Exposures were for 20, 40, and 60 min and included an intermittent exposure regimen (20 min on/20 min off), resulting in total incubation times of 20, 60, and 100 min, respectively. Concurrent controls were sham exposed. After extracting total cellular RNA, Northern blot analysis was used to assess the expression of the immediate early genes, c-fos and c-jun, in all cell populations. No change in c-fos transcript levels were detected after 20 min exposure at each field intensity (20 min was the only time period at which c-fos message could be detected consistently). Transcript levels for c-jun were altered only after 20 min exposure to 9 mW/cm² (average 38% decrease).

Jarupat S, Kawabata A, Tokura H, Borkiewicz A. Effects of the 1900 MHz Electromagnetic Field Emitted from Cellular Phone on Nocturnal Melatonin Secretion. *J Physiol Anthropol Appl Human Sci* 22(1):61-63, 2003.

Exposure to cellular phone EMF caused a significant reduction in salivary melatonin in female human subjects.

Jech R, Sonka K, Ruzicka E, Nebuzelsky A, Bohm J, Juklickova M, Nevsimalova S. Electromagnetic field of mobile phones affects visual event related potential in patients with narcolepsy. *Bioelectromagnetics* 22(7):519-528, 2001. The effects of the mobile phone (MP) electromagnetic fields on electroencephalography (EEG) and event-related potentials (ERP) were

examined. With regard to the reported effects of MP on sleep, 22 patients with narcolepsy-cataplexy were exposed or sham exposed for 45 min to the MP (900 MHz, specific absorption rate 0.06 W/kg) placed close to the right ear in a double blind study. There were no changes of the EEG recorded after the MP exposure. A subgroup of 17 patients was studied on visual ERP recorded during the MP exposure. Using an adapted "odd-ball" paradigm, each patient was instructed to strike a key whenever rare target stimuli were presented. There were three variants of target stimuli (horizontal stripes in (i) left, (ii) right hemifields or (iii) whole field of the screen). The exposure enhanced the positivity of the ERP endogenous complex solely in response to target stimuli in the right hemifield of the screen ($P < 0.01$). The reaction time was shortened by 20 ms in response to all target stimuli ($P < 0.05$). In conclusion, the electromagnetic field of MP may suppress the excessive sleepiness and improve performance while solving a monotonous cognitive task requiring sustained attention and vigilance.

Jensh RP, Behavioral teratologic studies using microwave radiation: is there an increased risk from exposure to cellular phones and microwave ovens? *Reprod Toxicol* 11(4):601-611, 1997. The objective of the investigations presented in this review was to determine if there are adverse effects due to chronic prenatal microwave exposure in rats at term and/or alterations in neonatal and adult offspring psychophysiologic development and growth. Following the establishment of a nonhyperthermal power density level of microwave radiation, pregnant rats were exposed throughout pregnancy to continuous wave 915 MHz, 2450 MHz, or 6000 MHz radiation at power density levels of 10, 20, or 35 mW/cm², respectively. Teratologic evaluation included the following parameters: maternal weight and weight gain; mean litter size; maternal organ weight and organ weight/body weight ratios; body weight ratios of brain, liver, kidneys, and ovaries; maternal peripheral blood parameters including hematocrit, hemoglobin, and white cell counts; number of resorptions and resorption rate; number of abnormalities and abnormality rate; mean term fetal weight. Mothers were rebred, and the second, nonexposed litters were evaluated for teratogenic effects. Exposed offspring were evaluated using the following perinatal and adult tests: eye opening, surface righting, negative geotaxis, auditory startle, air righting, open field, activity wheel, swimming, and forelimb hanging. Offspring were also monitored for weekly weight and weight gain. Animals exposed to 915 MHz did not exhibit any consistent significant alterations in any of the above parameters. Exposure to 2450 MHz resulted only in a significantly increased adult offspring activity level compared to nonexposed offspring. Offspring exposed to 6000 MHz radiation exhibited an initial slight, but significant, retardation in term weight, while mothers had a significantly reduced monocyte count. No changes in any of the other term parameters were observed. A few postnatal parameters were affected in offspring exposed to 6000 MHz. Weekly weights were lower in the exposed offspring, but they recovered by the fifth week. Eye opening was delayed, and there were changes in the water T-maze and open field performance levels. Several organ/body weight ratios differed from those of the control offspring. These results indicate that exposure

to 6000 MHz radiation at this power density level may result in subtle long-term neurophysiologic alterations. However, in the absence of a hyperthermic state, the microwave frequencies tested, which included frequencies used in cellular phones and microwave ovens, do not induce a consistent, significant increase in reproductive risk as assessed by classical morphologic and postnatal psychophysiologic parameters.

Johansen C, Boice JD, McLaughlin JK, Olsen JH, Cellular Telephones and Cancer-a Nationwide Cohort Study in Denmark. *J Natl Cancer Inst* 93(3):203-207, 2001. BACKGROUND: Use of cellular telephones is increasing exponentially and has become part of everyday life. Concerns about possible carcinogenic effects of radiofrequency signals have been raised, although they are based on limited scientific evidence. METHODS: A retrospective cohort study of cancer incidence was conducted in Denmark of all users of cellular telephones during the period from 1982 through 1995. Subscriber lists from the two Danish operating companies identified 420 095 cellular telephone users. Cancer incidence was determined by linkage with the Danish Cancer Registry. All statistical tests are two-sided. RESULTS: Overall, 3391 cancers were observed with 3825 expected, yielding a significantly decreased standardized incidence ratio (SIR) of 0.89 (95% confidence interval [CI] = 0.86 to 0.92). A substantial proportion of this decreased risk was attributed to deficits of lung cancer and other smoking-related cancers. No excesses were observed for cancers of the brain or nervous system (SIR = 0.95; 95% CI = 0.81 to 1.12) or of the salivary gland (SIR = 0.72; 95% CI = 0.29 to 1.49) or for leukemia (SIR = 0.97; 95% CI = 0.78-1.21), cancers of a priori interest. Risk for these cancers also did not vary by duration of cellular telephone use, time since first subscription, age at first subscription, or type of cellular telephone (analogue or digital). Analysis of brain and nervous system tumors showed no statistically significant SIRs for any subtype or anatomic location. CONCLUSIONS: The results of this investigation, the first nationwide cancer incidence study of cellular phone users, do not support the hypothesis of an association between use of these telephones and tumors of the brain or salivary gland, leukemia, or other cancers.

Johansen C, Boice JD Jr, McLaughlin JK, Christensen HC, Olsen JH. Mobile phones and malignant melanoma of the eye. *Brit J Cancer* 86:348-349, 2002.

Recently a four-fold increase in the risk of malignant melanoma of the eye was associated with the use of radiofrequency transmitting devices, including mobile phones in Germany. We contrasted the incidence rates of this rare cancer with the number of mobile phone subscribers in Denmark. We observed no increasing trend in the incidence rate of melanoma, which was in sharp contrast to the exponentially increasing number of mobile phone subscribers starting in the early 1980s. Our study provides no support for an association between mobile phones and ocular melanoma.

Kellenyi, L, Thuroczy, G, Faludy, B, Lenard, L, Effects of mobile GSM radiotelephone exposure on the auditory brainstem response (ABR). *Neurobiology* 7:79-81, 1999.

A 15-min exposure to GSM phone radiation caused an increase in auditory brainstem response in the exposed side of human subjects. Subjects also showed a hearing deficiency in the high frequency range (20 dB hearing deficiency from 2 KHz to 10 KHz).

Khudnitskii, SS, Moshkarev, EA, Fomenko, TV, [On the evaluation of the influence of cellular phones on their users]. [Article in Russian] *Med Tr Prom Ekol* (9):20-24, 1999.

The authors studied influence of ultrahigh frequency radiation caused by cellular phones on functional state of central nervous, cardiovascular systems and local temperature changes in cellular phones users. The head area near the phone antenna appeared to be under the most intensive heating. Ultrahigh frequency radiation induces significant changes in local temperature and in physiologic parameters of central nervous and cardiovascular systems.

Kimata H. Enhancement of allergic skin wheal responses by microwave radiation from mobile phones in patients with atopic eczema/dermatitis syndrome. *Int Arch Allergy Immunol* 129(4):348-350, 2002. Microwave radiation from mobile phones enhanced skin wheal responses induced by house dust mite and Japanese cedar pollen while it had no effect on wheal responses induced by histamine in patients with atopic eczema/dermatitis syndrome (AEDS). Microwave radiation also increased plasma levels of substance P (SP) and vasoactive intestinal peptide (VIP) in patients with AEDS. These results indicate that microwave radiation from mobile phones may enhance allergen-induced wheal responses in association with the release of SP and VIP. This finding may be useful in elucidating the pathophysiology and treatment of AEDS.

Koivisto, M, Revonsuo, A, Krause, C, Haarala, C, Sillanmaki, L, Laine, M, Hamalainen, H, Effects of 902 MHz electromagnetic field emitted by cellular telephones on response times in humans. *Neuroreport* 11(2):413-415, 2000.

The present study examined possible influences of a 902 MHz electromagnetic field emitted by cellular telephones on cognitive functioning in 48 healthy humans. A battery of 12 reaction time tasks was performed twice by each participant in a counterbalanced order: once with and once without the exposure to the field. The results showed that the exposure to the electromagnetic field speeded up response times in simple reaction time and vigilance tasks and that the cognitive time needed in a mental arithmetics task was decreased. The results suggest that exposure to the electromagnetic field emitted by cellular telephones may have a facilitatory effect on brain functioning, especially in tasks requiring attention and manipulation of information in working memory.

Koivisto M, Krause CM, Revonsuo A, Laine M, Hamalainen H, The effects of electromagnetic field emitted by GSM phones on working memory. *Neuroreport* 11(8):1641-1643, 2000.

The influence of pulsed radiofrequency (RF) electromagnetic fields of digital GSM mobile phones on working memory in healthy subjects were studied. Memory load was varied from 0 to 3 items in an n-back task. Each subject was tested twice within a single session, with and without the RF exposure (902MHz, 217Hz). The RF field speeded up response times when the memory load was three items but no effects of RF were observed with lower loads. The results suggest that RF fields have a measurable effect on human cognitive performance and encourage further studies on the interactions of RF fields with brain function.

Koivisto M, Haarala C, Krause CM, Revonsuo A, Laine M, Hamalainen H, GSM phone signal does not produce subjective symptoms.

***Bioelectromagnetics* 22(3):212-215, 2001.** The influence of pulsed radiofrequency (RF) electromagnetic fields of digital GSM mobile phones (902 MHz, 217 Hz pulse modulation) on subjective symptoms or sensations in healthy subjects were studied in two single-blind experiments. The duration of the RF exposure was about 60 min in Experiment 1 and 30 min in Experiment 2. Each subject rated symptoms or sensations in the beginning of the experimental session and at the end of both the exposure and the nonexposure conditions. The symptoms rated were headache, dizziness, fatigue, itching or tingling of the skin, redness on the skin, and sensations of warmth on the skin. The results did not reveal any differences between exposure and non-exposure conditions, suggesting that a 30-60 min exposure to this RF field does not produce subjective symptoms in humans.

Krause CM, Sillanmaki L, Koivisto M, Haggqvist A, Saarela C, Revonsuo A, Laine M, Hamalainen H, Effects of electromagnetic field emitted by cellular phones on the EEG during a memory task. *Neuroreport* 11(4):761-764, 2000.

The effects of electromagnetic fields (EMF) emitted by cellular phones on the ERD/ERS of the 4-6 Hz, 6-8 Hz, 8-10 Hz and 10-12 Hz EEG frequency bands were studied in 16 normal subjects performing an auditory memory task. All subjects performed the memory task both with and without exposure to a digital 902 MHz EMF in counterbalanced order. The exposure to EMF significantly increased EEG power in the 8-10 Hz frequency band only. Nonetheless, the presence of EMF altered the ERD/ERS responses in all studied frequency bands as a function of time and memory task (encoding vs retrieval). Our results suggest that the exposure to EMF does not alter the resting EEG per se but modifies the brain responses significantly during a memory task.

Krause CM, Sillanmaki L, Koivisto M, Haggqvist A, Saarela C, Revonsuo A, Laine M, Hamalainen H, Effects of electromagnetic fields emitted by cellular phones on the electroencephalogram during a visual working memory task. *Int J Radiat Biol* 76(12):1659-1667, 2000. PURPOSE: To examine the effects of electromagnetic fields (EMF) emitted by cellular phones on the event-related desynchronization/synchronization (ERD/ERS) responses of the 4-6, 6-8, 8-10 and 10-12Hz EEG frequency bands during cognitive processing. MATERIALS AND METHODS: Twenty-four subjects performed a visual sequential letter task (n-back task) with three different working memory load conditions: zero, one and two items. All subjects performed the memory task both with and without exposure to a digital 902 MHz EMF in counterbalanced order. RESULTS: The presence of EMF altered the ERD/ERS responses in the 6-8 and 8-10 Hz frequency bands but only when examined as a function of memory load and depending also on whether the presented stimulus was a target or not. CONCLUSIONS: The results suggest that the exposure to EMF modulates the responses of EEG oscillatory activity approximately 8 Hz specifically during cognitive processes.

Kwee S, Raskmark P, Changes in cell proliferation due to environmental non-ionizing radiation 2. Microwave radiation. *Bioelectrochem Bioenerg* 44(2) 251-255, 1998.

Due to the use of mobile telephones, there is an increased exposure of the environment to weak radiofrequency (RF) electromagnetic fields, emitted by these devices. This study was undertaken to investigate if the microwave radiation from these fields will have a similar effect on cell proliferation as weak electromagnetic (ELF) fields. The field was generated by signal simulation of the Global System for Mobile communications (GSM) of 960 MHz. Cell cultures, growing in microtiter plates, were exposed in a specially constructed chamber, a Transverse Electromagnetic (TEM) cell. The Specific Absorption Rate (SAR) values for each cell well were calculated for this exposure system. Experiments were performed on cell cultures of transformed human epithelial amnion cells (AMA), which were exposed to 960 MHz microwave fields at three different power levels and three different exposure times, respectively. It was found that cell growth in the exposed cells was decreased in comparison to that in the control and sham exposed cells. Cell proliferation during the period following exposure varied not only with the various SAR levels, but also with the length of exposure time. On the other hand, repeated periods of exposure did not seem to change the effects. There was a general linear correlation between power level and growth change. However, the exposure time required to obtain the maximum effect was not the same for the various power levels. It turned out that at low power level, a maximum effect was first reached after a longer exposure time than at higher power level. A similar phenomenon was registered in the studies on ELF electromagnetic fields. Here, it was found that there was a linear correlation between the length of exposure time to obtain maximum effect and field strength.

Kwee S, Raskmark P, Velizarov P. Changes in cellular proteins due to environmental non-ionizing radiation. I. Heat-shock proteins. *Electro- and Magnetobiology* 20: 141-152, 2001.

This paper describes the effect of weak microwave fields on the amounts of heat-shock proteins in cell cultures at various temperatures. The field was generated by signal simulation of the Global System for Mobile communications (GSM) of 960 Mhz, used in portable phones. Transformed human epithelial amnion (AMA) cells, growing on glass coverslips, were exposed in a transverse electromagnetic (TEM) cell to a microwave field, generating a specific absorption rate (SAR) of 2.1 mW.kg⁻¹ in the cells. Exposure temperatures were 35, 37, and 40 Å 0.1% C, respectively, and the exposure time was 20

min. The heat-shock proteins Hsp-70 and Hsp-27 were detected by immunofluorescence. Higher amounts of Hsp-70 were present in the cells exposed at 35 and 37°C than in the sham-exposed cells. These effects can be considered to be athermal, since the field strength was much lower than the safety standard for absence of heat generation by microwave fields. There was no significant response in the case of Hsp-27.

Lebedeva NN, Sulimov AV, Sulimova OP, Kotrovskaya TI, Gailus T, Cellular phone electromagnetic field effects on bioelectric activity of human brain. *Crit Rev Biomed Eng* 28(1-2):323-337, 2000. 24 volunteers participated in the experiments. The investigation of EEG reactions to cellular phone (EMF frequency 902.4 MHz and intensity 0.06 mW/cm²) was conducted. Two experiments were performed with each subject--cellular phone exposure and Placebo Duration of the experiment was 60 min: 15 min--background; 15 min--EMF exposure or Placebo; 30 min--afterexposure. EEG was recorded in 16 standard leads with "eyes open" and "eyes closed". Special software with non-linear dynamics was developed for EEG analyses. One parameter, multichannel (global) correlation dimension, was calculated. The changes of these parameters can be evidence of brain functional state changes. As a result of EEG record processing, a significant increase of global correlation dimension during the exposure and afterexposure period was discovered, more pronounced in the case of "eyes closed". That can be viewed as the manifestation of cortex activation under phone EMF exposure.

Lebedeva NN, Sulimov AV, Sulimova OP, Korotkovskaya TI, Gailus T, Investigation of brain potentials in sleeping humans exposed to the electromagnetic field of mobile phones. *Crit Rev Biomed Eng* 29(1):125-133, 2001. An investigation was made of 8-hour EEG tracings of sleeping humans exposed to the electromagnetic field of a GSM-standard mobile phone. To analyze the EEG-patterns, manual scoring, nonlinear dynamics, and spectral analysis were employed. It was found that, when human beings were exposed to the electromagnetic field of a cellular phone, their cerebral cortex biopotentials revealed an increase in the alpha-range power density as compared to the placebo experiment. It was also found that the dimension of EEG correlation dynamics and the relation of sleep stages changed under the influence of the electromagnetic field of a mobile phone.

Lee TMC, Ho SMY, Tsang LYH, Yang SYC, Li LSW, Chan CCH, Effect on human attention of exposure to the electromagnetic field emitted by mobile phones. *NeuroReport* 12:729-731, 2001.

This study examined the effect of exposure to the electromagnetic field emitted by mobile phones on human attention. Three measures of attention were administered to 72 teenagers, 37 of whom were mobile phone users. The results showed that the mobile phone users performed better on one of the three measures of attention than did the non-mobile phone users. The results suggest that exposure to the electromagnetic field emitted by mobile phones may have a mild facilitating effect on attention functions, which is consistent with previous

observations that exposure to the electromagnetic field has a facilitating effect on cognitive processing. The possibility that mobile phone users may be naturally better at multiple tasking tasks was discussed.

Leszczynski D, Joenväärä S, Reivinen J, Kuokka R, Non-thermal activation of the hsp27/p38MAPK stress pathway by mobile phone radiation in human endothelial cells: Molecular mechanism for cancer- and blood-brain barrier-related effects. *Differentiation* 70:120 – 129, 2002.

We have examined whether non-thermal exposures of cultures of the human endothelial cell line EA.hy926 to 900 MHz GSM mobile phone microwave radiation could activate stress response. Results obtained demonstrate that 1-hour non-thermal exposure of EA.hy926 cells changes the phosphorylation status of numerous, yet largely unidentified, proteins. One of the affected proteins was identified as heat shock protein-27 (hsp27). Mobile phone exposure caused a transient increase in phosphorylation of hsp27, an effect which was prevented by SB203580, a specific inhibitor of p38 mitogen-activated protein kinase (p38MAPK). Also, mobile phone exposure caused transient changes in the protein expression levels of hsp27 and p38MAPK. All these changes were non-thermal effects because, as determined using temperature probes, irradiation did not alter the temperature of cell cultures, which remained throughout the irradiation period at 37±0.3°C. Changes in the overall pattern of protein phosphorylation suggest that mobile phone radiation activates a variety of cellular signal transduction pathways, among them the hsp27/p38MAPK stress response pathway. Based on the known functions of hsp27, we put forward the hypothesis that mobile phone radiation-induced activation of hsp27 may (i) facilitate the development of brain cancer by inhibiting the cytochrome c/caspase-3 apoptotic pathway and (ii) cause an increase in blood-brain barrier permeability through stabilization of endothelial cell stress fibers. We postulate that these events, when occurring repeatedly over a long period of time, might become a health hazard because of the possible accumulation of brain tissue damage. Furthermore, our hypothesis suggests that other brain damaging factors may co-participate in mobile phone radiation-induced effects.

Li, JR, Chou, CK, McDougall, JA, Dasgupta, G, Wu, HH, Ren, RL, Lee, A, Han, J, Momand J TP53 tumor suppressor protein in normal human fibroblasts does not respond to 837 MHz microwave exposure. *Radiat Res* 151(6):710-716, 1999.

The TP53 tumor suppressor protein (formerly known as p53) responds to a wide variety of environmental insults. To evaluate the safety of cellular telephones, TP53 responses in human fibroblast cells were studied after exposure to 837 MHz microwaves. Cells were exposed in a temperature-controlled transverse electromagnetic (TEM) chamber to a specific absorption rate (SAR) of 0.9 or 9.0 W/kg at 837 MHz continuous-wave (CW) microwave irradiation for 2 h. The TP53 protein levels were measured by Western blot at 2, 8, 24 and 48 h after treatment. The TP53 protein levels in microwave-treated cells, sham-treated

cells, and untreated cells remained unchanged relative to each other at all times tested (Fisher test and Student-Newman-Keuls test, $P > 0.05$). No morphological alterations were observed in microwave-treated cells compared to sham-treated cells. We conclude that TP53 protein expression levels in cultured human fibroblast cells do not change significantly during a 48-h period after exposure to 837 MHz continuous microwaves for 2 h at SAR levels of 0.9 or 9.0 W/kg.

Li L, Bisht KS, LaGroye I, Zhang P, Straube WL, Moros EG, Roti Roti JL. Measurement of DNA damage in mammalian cells exposed in vitro to radiofrequency fields at sars of 3-5 w/kg. *Radiat Res* 156:328-332, 2001.

In the present study, we determined whether exposure of mammalian cells to 3.2-5.1 W/kg specific absorption rate (SAR) radiofrequency fields could induce DNA damage in murine C3H 10T(1/2) fibroblasts. Cell cultures were exposed to 847.74 MHz code-division multiple access (CDMA) and 835.62 frequency-division multiple access (FDMA) modulated radiations in radial transmission line (RTL) irradiators in which the temperature was regulated to 37.0 ± 0.3 degrees C. Using the alkaline comet assay to measure DNA damage, we found no statistically significant differences in either comet moment or comet length between sham-exposed cells and those exposed for 2, 4 or 24 h to CDMA or FDMA radiations in either exponentially growing or plateau-phase cells. Further, a 4-h incubation after the 2-h exposure resulted in no significant changes in comet moment or comet length. Our results show that exposure of cultured C3H 10T(1/2) cells at 37 degrees C CDMA or FDMA at SAR values of up to 5.1 W/kg did not induce measurable DNA damage.

Linz, KW, von Westphalen, C, Streckert, J, Hansen, V, Meyer, R, Membrane potential and currents of isolated heart muscle cells exposed to pulsed radio frequency fields. *Bioelectromagnetics* 20(8):497-511, 1999.

The influence of radio frequency (RF) fields of 180, 900, and 1800 MHz on the membrane potential, action potential, L-type Ca^{2+} current and potassium currents of isolated ventricular myocytes was tested. The study is based on 90 guinea-pig myocytes and 20 rat myocytes. The fields were applied in rectangular waveguides (1800 MHz at 80, 480, 600, 720, or 880 mW/kg and 900 MHz, 250 mW/kg) or in a TEM-cell (180 MHz, 80 mW/kg and 900 MHz, 15 mW/kg). Fields of 1800 and 900 MHz were pulsed according to the GSM-standard of cellular phones. The specific absorption rates were determined from computer simulations of the electromagnetic fields inside the exposure devices by considering the structure of the physiological test arrangement. The electrical membrane parameters were measured by whole cell patch-clamp. None of the tested electrophysiological parameters was changed significantly by exposure to RF fields. Another physical stimulus, lowering the temperature from 36 degrees C to 24 degrees C, decreased the current amplitude almost 50% and shifted the voltage dependence of the steady state activation parameter $d(\infty)$ and inactivation parameter $f(\infty)$ of L-type Ca^{2+} current by about 5 mV.

However, at this lower temperature RF effects (900 MHz, 250 mW/kg; 1800 MHz, 480 mW/kg) on L-type Ca(2+) current were also not detected.

Litovitz TA, Krause D, Penafiel M, Elson EC, Mullins JM, The role of coherence time in the effect of microwaves on ornithine decarboxylase activity. *Bioelectromagnetics* 14(5):395-403, 1993. Previously, we demonstrated the requirements for a minimum coherence time of an applied, small amplitude (10 microT) ELF magnetic field if the field were to produce an enhancement of ornithine decarboxylase activity in L929 fibroblasts. Further investigation has revealed a remarkably similar coherence time phenomenon for enhancement of ornithine decarboxylase activity by amplitude-modulated 915 MHz microwaves of large amplitude (SAR 2.5 W/kg). Microwave fields modulated at 55, 60, or 65 Hz approximately doubled ornithine decarboxylase activity after 8 h. Switching modulation frequencies from 55 to 65 Hz at coherence times of 1.0 s or less abolished enhancement, while times of 10 s or longer provided full enhancement. Our results show that the microwave coherence effects are remarkably similar to those observed with ELF fields.

Litovitz, TA, Penafiel, LM, Farrel, JM, Krause, D, Meister, R, Mullins, JM Bioeffects induced by exposure to microwaves are mitigated by superposition of ELF noise. *Bioelectromagnetics* 18(6):422-430, 1997.

We have previously demonstrated that microwave fields, amplitude modulated (AM) by an extremely low-frequency (ELF) sine wave, can induce a nearly twofold enhancement in the activity of ornithine decarboxylase (ODC) in L929 cells at SAR levels of the order of 2.5 W/kg. Similar, although less pronounced, effects were also observed from exposure to a typical digital cellular phone test signal of the same power level, burst modulated at 50 Hz. We have also shown that ODC enhancement in L929 cells produced by exposure to ELF fields can be inhibited by superposition of ELF noise. In the present study, we explore the possibility that similar inhibition techniques can be used to suppress the microwave response. We concurrently exposed L929 cells to 60 Hz AM microwave fields or a 50 Hz burst-modulated DAMPS (Digital Advanced Mobile Phone System) digital cellular phone field at levels known to produce ODC enhancement, together with band-limited 30-100 Hz ELF noise with root mean square amplitude of up to 10 microT. All exposures were carried out for 8 h, which was previously found to yield the peak microwave response. In both cases, the ODC enhancement was found to decrease exponentially as a function of the noise root mean square amplitude. With 60 Hz AM microwaves, complete inhibition was obtained with noise levels at or above 2 microT. With the DAMPS digital cellular phone signal, complete inhibition occurred with noise levels at or above 5 microT. These results suggest a possible practical means to inhibit biological effects from exposure to both ELF and microwave fields.

Loscher W, Kas G, Extraordinary behavior disorders in cows in proximity to transmission stations. *Der Praktische Tierarz* 79:437-444, 1998.

(Article in German)

In addition to reduction of milk yield and increased health problems, behavioral abnormalities were observed over a period of two years in a herd of dairy cows maintained in close proximity to a TV and cell phone transmitting antenna. Evaluation of possible factors which could explain the abnormalities in the live stock did not disclose any factors other than the high-frequency electromagnetic fields. An experiment in which a cow with abnormal behavior was brought to a stable 20 km away from the antenna resulted in a complete normalization of the cow within five days, whereas symptoms returned when the cow was brought back to the stable nearby the antenna. In view of the previous described effects of electromagnetic fields, it might be possible that the observed abnormalities in cows are related to electromagnetic field exposure. (power densities measured 0.02-7 mW/m²).

Maes A, Collier M, Slaets D, Verschaeve L, 954 MHz microwaves enhance the mutagenic properties of mitomycin C. *Environ Mol Mutagen* 28(1):26-30, 1996. This paper focuses on the combined effects of microwaves from mobile communication frequencies and a chemical DNA damaging agent mitomycin C (MMC). The investigation was performed in vitro by exposing whole blood samples to a 954 MHz emitting antenna from a GSM (Global System for Mobile Communication) base station, followed by lymphocyte cultivation in the presence of MMC. A highly reproducible synergistic effect was observed as based on the frequencies of sister chromatid exchanges in metaphase figures.

Maes A, Collier M, Van Gorp U, Vandoninck S, Verschaeve L, Cytogenetic effects of 935.2-MHz (GSM) microwaves alone and in combination with mitomycin C. *Mutat Res* 393(1-2):151-156, 1997.

This paper focuses on the genetic effects of microwaves from mobile communication frequencies (935.2 MHz) alone and in combination with a chemical DNA-damaging agent (mitomycin C). Three cytogenetic endpoints were investigated after in vitro exposure of human whole blood cells. These endpoints were the 'classical' chromosome aberration test, the sister chromatid exchange test and the alkaline comet assay. No direct cytogenetic effect was found. The combined exposure of the cells to the radiofrequency fields followed by their cultivation in the presence of mitomycin C revealed a very weak effect when compared to cells exposed to mitomycin C alone.

Maes A, Collier M, Verschaeve L Cytogenetic effects of 900 MHz (GSM) microwaves on human lymphocytes. *Bioelectromagnetics* 22(2):91-96, 2001 The cytogenetic effects of 900 MHz radiofrequency fields were investigated with the chromosome aberration and sister chromatid exchange frequency methods. Three different modes of exposure (continuous, pseudo-random and dummy burst) were studied for different power outputs (0, 2, 8, 15, 25, 50 W). The

specific absorption rates varied between 0 and 10 W/kg. We investigated the possible effects of the 900 MHz radiation alone as well as of combined exposure to the chemical or physical mutagens mitomycin C and X-rays. Overall, no indication was found of a mutagenic, and/or co-mutagenic/synergistic effect of this kind of nonionizing radiation.

Malyapa RS, Ahern EW, Straube WL, Moros EG, Pickard WF, Roti Roti JL, Measurement of DNA damage after exposure to electromagnetic radiation in the cellular phone communication frequency band (835.62 and 847.74 MHz). *Radiat Res* 148(6):618-627, 1997.

Mouse C3H 10T1/2 fibroblasts and human glioblastoma U87MG cells were exposed to cellular phone communication frequency radiations to investigate whether such exposure produces DNA damage in in vitro cultures. Two types of frequency modulations were studied: frequency-modulated continuous-wave (FMCW), with a carrier frequency of 835.62 MHz, and code-division multiple-access (CDMA) centered on 847.74 MHz. Exponentially growing (U87MG and C3H 10T1/2 cells) and plateau-phase (C3H 10T1/2 cells) cultures were exposed to either FMCW or CDMA radiation for varying periods up to 24 h in specially designed radial transmission lines (RTLs) that provided relatively uniform exposure with a specific absorption rate (SAR) of 0.6 W/kg. Temperatures in the RTLs were monitored continuously and maintained at 37 +/- 0.3 degrees C. Sham exposure of cultures in an RTL (negative control) and ¹³⁷Cs gamma-irradiated samples (positive control) were included with every experiment. The alkaline comet assay as described by Olive et al. (Exp. Cell Res. 198, 259-269, 1992) was used to measure DNA damage. No significant differences were observed between the test group exposed to FMCW or CDMA radiation and the sham-treated negative controls. Our results indicate that exposure of cultured mammalian cells to cellular phone communication frequencies under these conditions at an SAR of 0.6 W/kg does not cause DNA damage as measured by the alkaline comet assay.

Mann, K, Roschke, J, Effects of pulsed high-frequency electromagnetic fields on human sleep. *Neuropsychobiology* 33(1):41-47, 1996.

In the present study we investigated the influence of pulsed high-frequency electromagnetic fields of digital mobile radio telephones on sleep in healthy humans. Besides a hypnotic effect with shortening of sleep onset latency, a REM suppressive effect with reduction of duration and percentage of REM sleep was found. Moreover, spectral analysis revealed qualitative alterations of the EEG signal during REM sleep with an increased spectral power density. Knowing the relevance of REM sleep for adequate information processing in the brain, especially concerning mnemonic functions and learning processes, the results emphasize the necessity to carry out further investigations on the interaction of this type of electromagnetic fields and the human organism.

Mann, K, Roschke, J, Connemann, B, Beta, H, No effects of pulsed high-frequency electromagnetic fields on heart rate variability during human sleep. *Neuropsychobiology*;38(4):251-256, 1998.

The influence of pulsed high-frequency electromagnetic fields emitted by digital mobile radio telephones on heart rate during sleep in healthy humans was investigated. Beside mean RR interval and total variability of RR intervals based on calculation of the standard deviation, heart rate variability was assessed in the frequency domain by spectral power analysis providing information about the balance between the two branches of the autonomic nervous system. For most

parameters, significant differences between different sleep stages were found. In particular, slow-wave sleep was characterized by a low ratio of low- and high-frequency components, indicating a predominance of the parasympathetic over the sympathetic tone. In contrast, during REM sleep the autonomic balance was shifted in favor of the sympathetic activity. For all heart rate parameters, no significant effects were detected under exposure to the field compared to placebo condition. Thus, under the given experimental conditions, autonomic control of heart rate was not affected by weak-pulsed high-frequency electromagnetic fields.

Mann, K, Wagner, P, Brunn, G, Hassan, F, Hiemke, C, Roschke, J, Effects of pulsed high-frequency electromagnetic fields on the neuroendocrine system. *Neuroendocrinology* 67(2):139-144, 1998.

The influence of pulsed high-frequency electromagnetic fields emitted from a circularly polarized antenna on the neuroendocrine system in healthy humans was investigated (900 MHz electromagnetic field, pulsed with 217 Hz, average power density 0.02 mW/cm²). Nocturnal hormone profiles of growth hormone (GH), cortisol, luteinizing hormone (LH) and melatonin were determined under polysomnographic control. An alteration in the hypothalamo-pituitary-adrenal axis activity was found with a slight, transient elevation in the cortisol serum level immediately after onset of field exposure which persisted for 1 h. For GH, LH and melatonin, no significant effects were found under exposure to the field compared to the placebo condition, regarding both total hormone production during the entire night and dynamic characteristics of the secretion pattern. Also the evaluation of the sleep EEG data revealed no significant alterations under field exposure, although there was a trend to an REM suppressive effect. The results indicate that weak high-frequency electromagnetic fields have no effects on nocturnal hormone secretion except for a slight elevation in cortisol production which is transient, pointing to an adaptation of the organism to the stimulus.

Marino AA, Nilsen E, Frilot C. Nonlinear changes in brain electrical activity due to cell phone radiation. *Bioelectromagnetics* 24(5):339-346, 2003. We studied the effect of an electromagnetic field from a cellular telephone on brain electrical activity, using a novel analytical method based on a nonlinear model. The electroencephalogram (EEG) from rabbits was embedded in phase space and local recurrence plots were calculated and quantified using recurrence quantitation analysis to permit statistical comparisons between filtered segments of exposed and control epochs from individual rabbits. When the rabbits were exposed to the radiation from a standard cellular telephone (800 MHz band, 600 mW maximum radiated power) under conditions that simulated normal human use, the EEG was significantly affected in nine of ten animals studied. The effect occurred beginning about 100 ms after initiation of application of the field and lasted approximately 300 ms. In each case, the fields increased the randomness in the EEG. A control procedure ruled out the possibility that the observations were a product of the method of analysis. No differences were found between

exposed and control epochs in any animal when the experiment was repeated after the rabbits had been sacrificed, indicating that absorption of radiation by the EEG electrodes could not account for the observed effect. No effect was seen when deposition of energy in the brain was minimized by repositioning the radiating antenna from the head to the chest, showing that the type of tissue that absorbed the energy determined the observed changes in the EEG. We conclude that, in normal use, the fields from a standard cellular telephone can alter brain function as a consequence of absorption of energy by the brain.

Marino C, Cristalli G, Galloni P, Pasqualetti P, Piscitelli M, Lovisolo GA , Effects of microwaves (900 MHz) on the cochlear receptor: exposure systems and preliminary results. *Radiat Environ Biophys* 39(2):131-136, 2000.

The purpose of this paper is to present the experimental device and the work in progress performed in search for objective organic correlation of damage to hearing, examining possible acoustic otofunctional effects on the cochlear epithelium of the rat due to exposure to microwaves (900 MHz). Two experiments using male Sprague-Dawley rats were carried out with a far-field exposure in a cubic chamber. No statistically significant evidence was obtained at both specific absorption rate (SAR) values. The exposure system and the diagnostic apparatus are extremely useful to investigate a potential effect on the auditory system: however, with the parameters applied in these experiments, no evidence was observed.

Mashevich M, Folkman D, Kesar A, Barbul A, Korenstein R, Jerby E, Avivi L, Exposure of human peripheral blood lymphocytes to electromagnetic fields associated with cellular phones leads to chromosomal instability. *Bioelectromagnetics* 24:82-90, 2003.

Whether exposure to radiation emitted from cellular phones poses a health hazard is at the focus of current debate. We have examined whether in vitro exposure of human peripheral blood lymphocytes (PBL) to continuous 830 MHz electromagnetic fields causes losses and gains of chromosomes (aneuploidy), a major **somatic mutation** leading to genomic instability and thereby to cancer. PBL were irradiated at different average absorption rates (SAR) in the range of 1.6-8.8 W/kg for 72 hr in an exposure system based on a parallel plate resonator at temperatures ranging from 34.5-37.5 °C. The averaged SAR and its distribution in the exposed tissue culture flask were determined by combining measurements and numerical analysis based on a finite element simulation code. A linear increase in chromosome 17 aneuploidy was observed as a function of the SAR value, demonstrating that this radiation has a genotoxic effect. The SAR dependent aneuploidy was accompanied by an abnormal mode of replication of the chromosome 17 region engaged in segregation (repetitive DNA arrays associated with the centromere), suggesting that epigenetic

alterations are involved in the SAR dependent genetic toxicity. Control experiments (i.e., without any RF radiation) carried out in the temperature range of 34.5-38.5 °C showed that elevated temperature is not associated with either the genetic or epigenetic alterations observed following RF radiation - the increased levels of aneuploidy and the modification in replication of the centromeric DNA arrays. These findings indicate that the genotoxic effect of the electromagnetic radiation is elicited via a non-thermal pathway. Moreover, the fact that aneuploidy is a phenomenon known to increase the risk for cancer, should be taken into consideration in future evaluation of exposure guidelines.

Mausset A, de Seze R, Montpeyroux F, Privat A. Effects of radiofrequency exposure on the GABAergic system in the rat cerebellum: clues from semi-quantitative immunohistochemistry. *Brain Res* 912(1):33-46, 2001. The widespread use of cellular phones raises the problem of interaction of electromagnetic fields with the central nervous system (CNS). In order to measure these effects on neurotransmitter content in the CNS, we developed a protocol of neurotransmitter detection based on immunohistochemistry and image analysis. Gamma-vinyl-GABA (GVG), an inhibitor of the GABA-transaminase was injected in rats to increase GABA concentration in the CNS. The cellular GABA contents were then revealed by immunohistochemistry and semi-quantified by image analysis thanks to three parameters: optical density (O.D.), staining area, and number of positive cells. The increase in cerebellar GABA content induced by GVG 1200 mg/kg was reflected in these three parameters in the molecular and the granular layers. Therefore, control of immunohistochemistry parameters, together with appropriate image analysis, allowed both the location and the detection of variations in cellular neurotransmitter content. This protocol was used to investigate the effects of exposure to 900 MHz radiofrequencies on cerebellar GABA content. Both pulsed emission with a specific absorption rate (SAR) of 4 W/kg and continuous emission with high SAR (32 W/kg) were tested. We observed a selective diminution of the stained processes area in the Purkinje cell layer after exposure to pulsed radiofrequency and, in addition, a decrease in O.D. in the three cell layers after exposure to continuous waves. Whether this effect is, at least partly, due to a local heating of the tissues is not known. Overall, it appears that high energetic radiofrequency exposure induces a diminution in cellular GABA content in the cerebellum.

McNamee JP, Bellier PV, Gajda GB, Miller SM, Lemay EP, Lavallee BF, Marro L, Thansandote A. DNA Damage and Micronucleus Induction in Human Leukocytes after Acute In Vitro Exposure to a 1.9 GHz Continuous-Wave Radiofrequency Field. *Radiat Res* 158(4):523-533, 2002a. Human blood cultures were exposed to a 1.9 GHz continuous-wave (CW) radiofrequency (RF) field for 2 h using a series of six circularly polarized, cylindrical waveguides. Mean specific absorption rates (SARs) of 0.0, 0.1, 0.26, 0.92, 2.4 and 10 W/kg were achieved, and the temperature within the cultures during a 2-h exposure

was maintained at 37.0 +/- 0.5 degrees C. Concurrent negative (incubator) and positive (1.5 Gy (137)Cs gamma radiation) control cultures were run for each experiment. DNA damage was quantified immediately after RF-field exposure using the alkaline comet assay, and four parameters (tail ratio, tail moment, comet length and tail length) were used to assess DNA damage for each comet. No evidence of increased primary DNA damage was detected by any parameter for RF-field-exposed cultures at any SAR tested. The formation of micronuclei in the RF-field-exposed blood cell cultures was assessed using the cytokinesis-block micronucleus assay. There was no significant difference in the binucleated cell frequency, incidence of micronucleated binucleated cells, or total incidence of micronuclei between any of the RF-field-exposed cultures and the sham-exposed controls at any SAR tested. These results do not support the hypothesis that acute, nonthermalizing 1.9 GHz CW RF-field exposure causes DNA damage in cultured human leukocytes.

McNamee JP, Bellier PV, Gajda GB, Lavallee BF, Lemay EP, Marro L, Thansandote A. DNA Damage in Human Leukocytes after Acute In Vitro Exposure to a 1.9 GHz Pulse-Modulated Radiofrequency Field. *Radiat Res* 158(4):534-537, 2002b. Blood cultures from human volunteers were exposed to an acute 1.9 GHz pulse-modulated radiofrequency (RF) field for 2 h using a series of six circularly polarized, cylindrical waveguides. Mean specific absorption rates (SARs) ranged from 0 to 10 W/kg, and the temperature within the cultures during the exposure was maintained at 37.0 +/- 0.5 degrees C. DNA damage was quantified in leukocytes by the alkaline comet assay and the cytokinesis-block micronucleus assay. When compared to the sham-treated controls, no evidence of increased primary DNA damage was detected by any parameter for any of the RF-field-exposed cultures when evaluated using the alkaline comet assay. Furthermore, no significant differences in the frequency of binucleated cells, incidence of micronucleated binucleated cells, or total incidence of micronuclei were detected between any of the RF-field-exposed cultures and the sham-treated control at any SAR tested. These results do not support the hypothesis that acute, nonthermalizing 1.9 GHz pulse-modulated RF-field exposure causes DNA damage in cultured human leukocytes.

McNamee JP, Bellier PV, Gajda GB, Lavallee BF, Marro L, Lemay E, Thansandote A. No Evidence for Genotoxic Effects from 24 h Exposure of Human Leukocytes to 1.9 GHz Radiofrequency Fields. *Radiat Res* 159(5):693-697, 2003. McNamee, J. P., Bellier, P. V., Gajda, G. B., Lavallee, B. F., Marro, L., Lemay, E. and Thansandote, A. No Evidence for Genotoxic Effects from 24 h Exposure of Human Leukocytes to 1.9 GHz Radiofrequency Fields. *Radiat. Res.* 159, 693-697 (2003). The current study extends our previous investigations of 2-h radiofrequency (RF)-field exposures on genotoxicity in human blood cell cultures by examining the effect of 24-h continuous-wave (CW) and pulsed-wave (PW) 1.9 GHz RF-field exposures on both primary DNA damage and micronucleus induction in human leukocyte cultures. Mean specific absorption rates (SARs) ranged from 0 to 10 W/kg, and the temperature within the cultures was maintained at 37.0 +/- 1.0 degrees C for the duration of the 24-h

exposure period. No significant differences in primary DNA damage were observed between the sham-treated controls and any of the CW or PW 1.9 GHz RF-field-exposed cultures when processed immediately after the exposure period by the alkaline comet assay. Similarly, no significant differences were observed in the incidence of micronuclei, incidence of micronucleated binucleated cells, frequency of binucleated cells, or proliferation index between the sham-treated controls and any of the CW or PW 1.9 GHz RF-field-exposed cultures. In conclusion, the current study found no evidence of 1.9 GHz RF-field-induced genotoxicity in human blood cell cultures after a 24-h exposure period.

Morrissey JJ, Raney S, Heasley E, Rathinavelu P, Dauphinee M, Fallon JH, IRIDIUM exposure increases c-fos expression in the mouse brain only at levels which likely result in tissue heating. *Neuroscience* 92(4):1539-1546, 1999.

With the rapid development of wireless communication technology over the last 20 years, there has been some public concern over possible health effects of long-term, low-level radiofrequency exposure from cellular telephones. As an initial step in compiling a database for risk analysis by government agencies, the effects of 1-h exposure of mice to a 1.6-GHz radiofrequency signal, given as either a continuous wave or pulse modulated at 11 Hz with a duty cycle of 4:1 and a pulse duration of 9.2 ms (IRIDIUM), on c-fos gene expression in the brain was investigated. The IRIDIUM signal is the operating frequency for a ground-to-satellite-to-ground cellular communications web which has recently become fully operational, and was named as such due to the original designed employment of the same number of low orbiting satellites as there are electrons orbiting the nucleus of an iridium atom. The expression of c-fos was not significantly elevated in the brains of mice until exposure levels exceeded six times the peak dose and 30 times the whole body average dose as maximal cellular telephone exposure limits in humans. Higher level exposure using either continuous wave (analog) or IRIDIUM signals elevated c-fos to a similar extent, suggesting no obvious pulsed modulation-specific effects. The pattern of c-fos elevation in limbic cortex and subcortex areas at higher exposure levels is most consistent with a stress response due to thermal perception coupled with restraint and/or neuron activity near thermoregulatory regions, and not consistent with any direct interaction of IRIDIUM energy with brain tissue.

Moustafa YM, Moustafa RM, Belacy A, Abou-El-Ela SH, Ali FM. Effects of acute exposure to the radiofrequency fields of cellular phones on plasma lipid peroxide and antioxidase activities in human erythrocytes. *J Pharm Biomed Anal* 26(4):605-608, 2001. Radiofrequency fields of cellular phones may affect biological systems by increasing free radicals, which appear mainly to enhance lipid peroxidation, and by changing the antioxidase activities of human blood thus leading to oxidative stress. To test this, we have investigated the effect of acute exposure to radiofrequency fields of commercially available

cellular phones on some parameters indicative of oxidative stress in 12 healthy adult male volunteers. Each volunteer put the phone in his pocket in standby position with the keypad facing the body. The parameters measured were lipid peroxide and the activities of superoxide dismutase (SOD), total glutathione peroxidase (GSH-Px) and catalase. The results obtained showed that the plasma level of lipid peroxide was significantly increased after 1, 2 and 4 h of exposure to radiofrequency fields of the cellular phone in standby position. Moreover, the activities of SOD and GSH-Px in human erythrocytes showed significant reduction while the activity of catalase in human erythrocytes did not decrease significantly. These results indicate that acute exposure to radiofrequency fields of commercially available cellular phones may modulate the oxidative stress of free radicals by enhancing lipid peroxidation and reducing the activation of SOD and GSH-Px, which are free radical scavengers. Therefore, these results support the interaction of radiofrequency fields of cellular phones with biological systems.

Muscat JE, Malkin MG, Thompson S, Shore RE, Stellman SD, McRee D, Neugut AI, Wynder EL, Handheld cellular telephone use and risk of brain cancer. JAMA 284(23):3001-3007, 2000. CONTEXT: A relative paucity of data exist on the possible health effects of using cellular telephones. OBJECTIVE: To test the hypothesis that using handheld cellular telephones is related to the risk of primary brain cancer. DESIGN AND SETTING: Case-control study conducted in 5 US academic medical centers between 1994 and 1998 using a structured questionnaire. PATIENTS: A total of 469 men and women aged 18 to 80 years with primary brain cancer and 422 matched controls without brain cancer. MAIN OUTCOME MEASURE: Risk of brain cancer compared by use of handheld cellular telephones, in hours per month and years of use. RESULTS: The median monthly hours of use were 2.5 for cases and 2.2 for controls. Compared with patients who never used handheld cellular telephones, the multivariate odds ratio (OR) associated with regular past or current use was 0.85 (95% confidence interval [CI], 0.6-1.2). The OR for infrequent users (<0.72 h/mo) was 1.0 (95% CI, 0.5-2.0) and for frequent users (>10.1 h/mo) was 0.7 (95% CI, 0.3-1.4). The mean duration of use was 2.8 years for cases and 2.7 years for controls; no association with brain cancer was observed according to duration of use ($P = .54$). In cases, cerebral tumors occurred more frequently on the same side of the head where cellular telephones had been used (26 vs 15 cases; $P = .06$), but in the cases with temporal lobe cancer a greater proportion of tumors occurred in the contralateral than ipsilateral side (9 vs 5 cases; $P = .33$). The OR was less than 1.0 for all histologic categories of brain cancer except for uncommon neuroepitheliomatous cancers (OR, 2.1; 95% CI, 0.9-4.7). CONCLUSIONS: Our data suggest that use of handheld cellular telephones is not associated with risk of brain cancer, but further studies are needed to account for longer induction periods, especially for slow-growing tumors with neuronal features.

Muscat JE, Malkin MG, Shore RE, Thompson S, Neugut AL, Stellman SD, Bruce J. Handheld cellular telephones and risk of acoustic neuroma. *Neurology* 58:1304-1306, 2002.

The hypothesis that intracranial energy deposition from handheld cellular telephones causes acoustic neuroma was tested in an epidemiologic study of 90 patients and 86 control subjects.

The relative risk was 0.9 ($p = 0.07$) and did not vary significantly by the frequency, duration, and lifetime hours of use. In patients who used cellular telephones, the tumor occurred more often on the contralateral than ipsilateral side of the head. Further efforts should focus on potentially longer induction periods.

Nakamura H, Matsuzaki I, Hatta K, Nobukuni Y, Kambayashi Y, Ogino K. Nonthermal effects of mobile-phone frequency microwaves on uteroplacental functions in pregnant rats. *Reprod Toxicol* 2003 17(3):321-326, 2003. Exposure to high-density microwaves can cause detrimental effects on the testis, eye, and other tissues, and induce significant biologic changes through thermal actions. To examine nonthermal effect of continuous wave (CW) 915MHz microwaves used in cellular phones, we compared the effects of microwaves with those of heat. Thirty-six pregnant rats were assigned to six groups: rats exposed to microwaves at 0.6 or 3mW/cm(2) incident power density at 915MHz for 90min, rats immersed in water at 38 or 40 degrees C, which induces about the same increase in colonic temperature of 1.0 or 3.5 degrees C as 0.6 or 3mW/cm(2) microwaves, respectively; rats immersed in water at 34 degrees C, which is considered to be thermoneutral; and control rats. We identified significant differences in the uteroplacental circulation, and in placental endocrine and immune functions between pregnant rats immersed in water at 34 and 38 degrees C, but not between rats immersed at 38 degrees C and those exposed to microwaves at 0.6mW/cm(2). By contrast, we observed significant decreases in uteroplacental blood flow and estradiol in rats exposed to microwaves at 3mW/cm(2) as compared with those immersed in water at 40 degrees C. These results suggest microwaves at 0.6mW/cm(2) at 915MHz, equal to a specific absorption rate (SAR) of 0.4W/kg, which is the maximum permissible exposure level recommended by the American National Standards Institute (ANSI), do not exert nonthermal effects on blood estradiol and progesterone, on splenic natural killer cell activity, on the uteroplacental circulation.

Oftedal G, Wilen J, Sandstrom M, Mild KH, Symptoms experienced in connection with mobile phone use. *Occup Med (Lond)* 50(4):237-245, 2000.

Many people in Norway and Sweden reported headaches, fatigue, and other symptoms experienced in connection with the use of a mobile phone (MP). Therefore, we initiated a cross-sectional epidemiological study among 17,000 people, all using an MP in their job. Thirty-one percent of the respondents in Norway and 13% of those in Sweden had experienced at least one symptom in connection with MP use. Next to the sensations of warmth on the ear and behind/around the ear, burning sensations in the facial skin and headaches were most commonly reported. Most symptoms usually began during or within half an hour after the call and lasted for up to 2 h. Relatively few had consulted a physician or been on sick leave because of the symptoms, but about 45% among those with an MP attributed symptom had taken steps to reduce the symptom. These results suggest an awareness of the symptoms, but not necessarily a serious health problem.

Pacini S, Ruggiero M, Sardi I, Aterini S, Gulisano F, Gulisano M. Exposure to global system for mobile communication (GSM) cellular phone radiofrequency alters gene expression, proliferation, and morphology of human skin fibroblasts. *Oncol Res* 13(1):19-24, 2002.

Human skin fibroblasts were exposed to global system for mobile communication (GSM) cellular phone radiofrequency for 1 h. GSM exposure induced alterations in cell morphology and increased the expression of mitogenic signal transduction genes (e.g., MAP kinase kinase 3, G2/mitotic-specific cyclin G1), cell growth inhibitors (e.g., transforming growth factor-beta), and genes controlling apoptosis (e.g., bax). A significant increase in DNA synthesis and intracellular mitogenic second messenger formation matched the high expression of MAP kinase family genes. These findings show that these electromagnetic fields have significant biological effects on human skin fibroblasts.

Paredi P, Kharitonov SA, Hanazawa T, Barnes PJ, Local vasodilator response to mobile phones. *Laryngoscope* 111(1):159-162, 2001.

OBJECTIVES: The use of mobile phones with the resulting generation of potentially harmful electromagnetic fields (EMF) is the focus of public interest. Heat generation and the activation of the inducible form of nitric oxide (NO) synthase may be possible causes of the biological effects of EMF exposure. We investigated if a mobile telephone conversation can modify skin temperature, NO, and nasal resistance. **METHODS:** We studied the effect of an EMF (900 MHz) generated by a commercially available cellular phone during a 30-minute telephone conversation on skin temperature, nasal NO measured by chemiluminescence, and nasal minimal cross-sectional area (MCA) measured by rhinometry. Eleven normal subjects (mean age \pm standard error of mean [SEM], 32 \pm 5 y; 10 male) were studied. **RESULTS:** There was a similar and significant increase in skin temperature of the nostril and occipital area on the same side as the telephone (maximal increase 2.3 \pm 0.2 degrees C at 6 min) as well as a tendency for higher nasal NO levels (maximal increase 12.9 \pm 4.9% at 10 min), whereas the MCA was significantly reduced (maximal decrease -27 \pm 6% at 15 min). Such changes were not recorded when an earpiece was used to avoid the direct exposure to the electromagnetic field. There were no changes in the skin temperature and nasal NO measured on the opposite side to the mobile phone, whereas the MCA was significantly increased (38 \pm 10%). **CONCLUSIONS:** Exposure to EMF produced by a mobile phone produces biological effects that can be easily measured. Microwaves may increase skin temperature and therefore cause vasodilation and reduce MCA. Further studies are needed to study the long-term effects of mobile phone use and the relation among NO production, vasodilation, and temperature.

Penafiel LM, Litovitz T, Krause D, Desta A, Mullins JM, Role of modulation on the effect of microwaves on ornithine decarboxylase activity in L929 cells. *Bioelectromagnetics* 18(2):132-141, 1997.

The effect of 835 MHz microwaves on the activity of ornithine decarboxylase (ODC) in L929 murine cell was investigated at an SAR of approximately 2.5 W/kg. The results depended upon the type of modulation employed. AM frequencies of 16 Hz and 60 Hz produced a transient increase in ODC activity that reached a peak at 8 h of exposure and returned to control levels after 24 h of

exposure. In this case, ODC was increased by a maximum of 90% relative to control levels. A 40% increase in ODC activity was also observed after 8 h of exposure with a typical signal from a TDMA digital cellular telephone operating in the middle of its transmission frequency range (approximately 840 MHz). This signal was burst modulated at 50 Hz, with approximately 30% duty cycle. By contrast, 8 h exposure with 835 MHz microwaves amplitude modulated with speech produced no significant change in ODC activity. Further investigations, with 8 h of exposure to AM microwaves, as a function of modulation frequency, revealed that the response is frequency dependent, decreasing sharply at 6 Hz and 600 Hz. Exposure with 835 MHz microwaves, frequency modulated with a 60 Hz sinusoid, yielded no significant enhancement in ODC activity for exposure times ranging between 2 and 24 h. Similarly, exposure with a typical signal from an AMPS analog cellular telephone, which uses a form of frequency modulation, produced no significant enhancement in ODC activity. Exposure with 835 MHz continuous wave microwaves produced no effects for exposure times between 2 and 24 h, except for a small but statistically significant enhancement in ODC activity after 6 h of exposure. Comparison of these results suggests that effects are much more robust when the modulation causes low-frequency periodic changes in the amplitude of the microwave carrier.

Persson BRR, Salford LG, Brun A, Blood-brain barrier permeability in rats exposed to electromagnetic fields used in wireless communication. *Wireless Network* 3:455-461, 1997.

Biological effects of radio frequency electromagnetic fields (EMF) on the blood-brain barrier (BBB) have been studied in Fischer 344 rats of both sexes. The rats were not anesthetised during the exposure. The brains were perfused with saline for 3-4 minutes, and thereafter perfusion fixed with 4% formaldehyde for 5-6 minutes. Whole coronal sections of the brains were dehydrated and embedded in paraffin and sectioned at 5 micrometers. Albumin and fibrinogen were demonstrated immunochemically and classified as normal versus pathological leakage. In the present investigation we exposed male and female Fischer 344 rats in a Transverse Electromagnetic Transmission line chamber to microwaves of 915 MHz as continuous wave (CW) and pulse-modulated with different pulse power and at various time intervals. The CW-pulse power varied from 0.001 W to 10 W and the exposure time from 2 min to 960 min. In each experiment we exposed 4-6 rats with 2-4 controls randomly placed in excited and non-excited TEM cells, respectively. We have in total investigated 630 exposed rats at various modulation frequencies and 372 controls. The frequency of pathological rats is significantly increased ($P < 0.0001$) from 62/372 (ratio 0.17 ± 0.02) for control rats to 244/630 (ratio: 0.39 ± 0.043) in all exposed rats. Grouping the exposed animals according to the level or specific absorption energy (J/kg) give significant difference in all levels above 1.5 J/kg. The exposure was 915 MHz microwaves either pulse modulated (PW) at 217 Hz with 0.57 ms pulse width, at 50 Hz with 6.6 ms pulse width or continuous wave (CW). The frequency of pathological rats (0.17) among controls in the various groups is not significantly

different. The frequency of pathological rats was 170/480 (0.35 ± 0.03) among rats exposed to pulse modulated (PW) and 74/149 (0.50 ± 0.07) among rats exposed to continuous wave exposure (CW). These results are both highly significantly different to their corresponding controls ($p < 0.0001$) and the frequency of pathological rats after exposure to pulsed radiation (PW) is significantly less ($p < 0.002$) than after exposure to continuous wave radiation (CW).

Phillips, J.L., Ivaschuk, O., Ishida-Jones, T., Jones, R.A., Campbell-Beachler, M. and Haggren, W. DNA damage in Molt-4 T- lymphoblastoid cells exposed to cellular telephone radiofrequency fields in vitro. *Bioelectrochem. Bioenerg.* 45:103-110, 1998.

Molt-4 T-lymphoblastoid cells have been exposed to pulsed signals at cellular telephone frequencies of 813.5625 MHz (iDEN signal) and 836.55 MHz (TDMA signal). These studies were performed at low SAR (average = 2.4 and 24 microwatt/g for iDEN and 2.6 and 26 microwatt/g for TDMA) in studies designed to look for athermal RF effects. The alkaline comet, or single cell gel electrophoresis, assay was employed to measure DNA single-strand breaks in cell cultures exposed to the radiofrequency (RF) signal as compared to concurrent sham-exposed cultures. Tail moment and comet extent were calculated as indicators of DNA damage. Statistical differences in the distribution of values for tail moment and comet extent between exposed and control cell cultures were evaluated with the Kolmogorov-Smirnoff distribution test. Data points for all experiments of each exposure condition were pooled and analyzed as single groups. It was found that: 1) exposure of cells to the iDEN signal at an SAR of 2.4 microwatt/g for 2 h or 21 h significantly decreased DNA damage; 2) exposure of cells to the TDMA signal at an SAR of 2.6 microwatt/g for 2 h and 21 h significantly decreased DNA damage; 3) exposure of cells to the iDEN signal at an SAR of 24 microwatt/g for 2 h and 21 h significantly increased DNA damage; 4) exposure of cells to the TDMA signal at an SAR of 26 microwatt/g for 2 h significantly decreased DNA damage. The data indicate a need to study the effects of exposure to RF signals on direct DNA damage and on the rate at which DNA damage is repaired.

Philippova TM, Novoselov VI, Alekseev SI, Influence of microwaves on different types of receptors and the role of peroxidation of lipids on receptor-protein shedding. *Bioelectromagnetics* 15(3):183-192, 1994. The effects of a continuous wave or pulse-modulated, 900 MHz microwave field were studied by in vitro assays of rat chemoreceptors. The pulsed field was modulated as rectangular waves at rates of 1, 6, 16, 32, 75, or 100 pps. The pulse-period to pulse-duration ratio was 5 in all cases, and specific absorption rates (SARs) ranged from 0.5 to 18 W/kg. Binding of ligands to cell membranes was differentially affected by exposure to microwaves. For example, binding of H³-glutamic acid to hippocampal cells was not altered by a 15 min exposure to a continuous wave field at 1 W/kg, but binding of H³-dihydroalprenolol to liver-cell

membranes of neonates underwent a fivefold decrease under the same field conditions. This effect was not dependent on modulation or on a change in the constant of stimulus-receptor binding but depended on a shedding of the membrane's receptor elements into solution. The magnitude of inhibition correlated with the oxygen concentration in the exposed suspension. Antioxidants (dithiothreitol and ionol) inhibited the shedding of receptor elements. The microwave exposure did not cause an accumulation of products from the peroxidation of lipids (POL). Ascorbate-dependent or non-enzymatic POL was not responsible for the inhibition, and POL was not found in other model systems. However, enzymatic POL mechanisms in localized areas of receptor binding remain a possibility.

Preece, AW, Iwi, G, Davies-Smith, A, Wesnes, K, Butler, S, Lim, E, Varey, A, Effect of a 915-MHz simulated mobile phone signal on cognitive function in man. *Int J Radiat Biol* 75(4):447-456, 1999.

PURPOSE: To examine whether a simulated mobile telephone transmission at 915 MHz has an effect on cognitive function in man. **MATERIALS AND METHODS:** Thirty-six subjects in two groups were each given two training sessions and then three test sessions in a randomized three-way cross-over design. About 1 W mean power at 915 MHz from a quarter-wave antenna mounted on a physical copy of an analogue phone, as a sine wave, or modulated at 217 Hz with 12.5% duty cycle, or no power, was applied to the left squamous temple region of the subjects while they undertook a series of cognitive function tests lasting approximately 25-30 min. The second group was investigated for sleep, consumption of alcohol and beverages, and any other substances that might affect performance. **RESULTS:** In both groups, the only test affected was the choice reaction time and this showed as an increase in speed (a decrease in reaction time). There were no changes in word, number or picture recall, or in spatial memory. While an effect of visit-order was evident suggesting a learning effect of repeat tests, the design of the study allowed for this. Additionally, there was no systematic error introduced as a result of consumption of substances or sleep time. **CONCLUSIONS:** There was evidence of an increase in responsiveness, strongly in the analogue and less in the digital simulation, in choice reaction time. This could be associated with an effect on the angular gyrus that acts as an interface between the visual and speech centres and which lies directly under and on the same side as the antenna. Such an effect could be consistent with mild localized heating, or possibly a non-thermal response, which is nevertheless power-dependent.

Repacholi, MH, Basten, A, Gebiski, V, Noonan, D, Finnie, J, Harris, AW, Lymphomas in E mu-Pim1 transgenic mice exposed to pulsed 900 MHz electromagnetic fields. *Radiat Res* 147(5):631-640, 1997.

Whether radiofrequency (RF) fields are carcinogenic is controversial; epidemiological data have been inconclusive and animal tests limited. The aim of the present study was to determine whether long-term exposure to pulse-

modulated RF fields similar to those used in digital mobile telecommunications would increase the incidence of lymphoma in E mu-Pim1 transgenic mice, which are moderately predisposed to develop lymphoma spontaneously. One hundred female E mu-Pim1 mice were sham-exposed and 101 were exposed for two 30-min periods per day for up to 18 months to plane-wave fields of 900 MHz with a pulse repetition frequency of 217 Hz and a pulse width of 0.6 ms. Incident power densities were 2.6-13 W/m² and specific absorption rates were 0.008-4.2 W/kg, averaging 0.13-1.4 W/kg. Lymphoma risk was found to be significantly higher in the exposed mice than in the controls (OR = 2.4. P = 0.006, 95% CI = 1.3-4.5). Follicular lymphomas were the major contributor to the increased tumor incidence. Thus long-term intermittent exposure to RF fields can enhance the probability that mice carrying a lymphomagenic oncogene will develop lymphomas. We suggest that such genetically cancer-prone mice provide an experimental system for more detailed assessment of dose-response relationships for risk of cancer after RF-field exposure.

Roschke, J, Mann, K, No short-term effects of digital mobile radio telephone on the awake human electroencephalogram. *Bioelectromagnetics* 18(2):172-176, 1997.

A recent study reported the results of an exploratory study of alterations of the quantitative sleep profile due to the effects of a digital mobile radio telephone. Rapid eye movement (REM) was suppressed, and the spectral power density in the 8-13 Hz frequency range during REM sleep was altered. The aim of the present study was to illuminate the influence of digital mobile radio telephone on the awake electroencephalogram (EEG) of healthy subjects. For this purpose, we investigated 34 male subjects in a single-blind cross-over design experiment by measuring spontaneous EEGs under closed-eyes condition from scalp positions C3 and C4 and comparing the effects of an active (0.05 mW/cm²) and an inactive digital mobile radio telephone (GSM) system. During exposure of nearly 3.5 min to the 900 MHz electromagnetic field pulsed at a frequency of 217 Hz and with a pulse width of 580 microseconds, we could not detect any difference in the awake EEGs in terms of spectral power density measures.

Roti Roti JL , Malyapa RS, Bisht KS, Ahern EW, Moros EG, Pickard WF, Straube WL, Neoplastic Transformation in C3H 10T(1/2) Cells after Exposure to 835.62 MHz FDMA and 847.74 MHz CDMA Radiations. *Radiat Res* 155(1):239-247, 2001. Roti Roti, J. L., Malyapa, R. S., Bisht, K. S., Ahern, E. W., Moros, E. G., Pickard, W. F. and Straube, W. L. Neoplastic Transformation in C3H 10T(1/2) Cells after Exposure to 835.62 MHz FDMA and 847.74 MHz CDMA Radiations. The effect of radiofrequency (RF) radiation in the cellular phone communication range (835.62 MHz frequency division multiple access, FDMA; 847.74 MHz code division multiple access, CDMA) on neoplastic transformation frequency was measured using the in vitro C3H 10T(1/2) cell transformation assay system. To determine if 835.62 MHz FDMA or 847.74 MHz CDMA radiations have any genotoxic effects that induce neoplastic

transformation, C3H 10T(1/2) cells were exposed at 37 degrees C to either of the above radiations [each at a specific absorption rate (SAR) of 0.6 W/kg] or sham-exposed at the same time for 7 days. After the culture medium was changed, the cultures were transferred to incubators and refed with fresh growth medium every 7 days. After 42 days, the cells were fixed and stained with Giemsa, and transformed foci were scored. To determine if exposure to 835.62 MHz FDMA or 847.74 MHz CDMA radiation has any epigenetic effects that can promote neoplastic transformation, cells were first exposed to 4.5 Gy of X rays to induce the transformation process and then exposed to the above radiations (SAR = 0.6 W/kg) in temperature-controlled irradiators with weekly refeeding for 42 days. After both the 7-day RF exposure and the 42-day RF exposure after X irradiation, no statistically significant differences in the transformation frequencies were observed between incubator controls, the sham-exposed (maintained in irradiators without power to the antenna), and the 835.62 MHz FDMA or 847.74 MHz CDMA-exposed groups

Salford LG, Brun A, Stureson K, Eberhardt JL, Persson BR Permeability of the blood-brain barrier induced by 915 MHz electromagnetic radiation, continuous wave and modulated at 8, 16, 50, and 200 Hz. *Microsc Res Tech* 27(6):535-542, 1994. Biological effects of electromagnetic fields (EMF) on the blood-brain barrier (BBB) can be studied in sensitive and specific models. In a previous investigation of the permeability of the blood-brain barrier after exposure to the various EMF-components of proton magnetic resonance imaging (MRI), we found that the exposure to MRI induced leakage of Evans Blue labeled proteins normally not passing the BBB of rats [Salford et al. (1992), in: *Resonance Phenomena in Biology*, Oxford University Press, pp. 87-91]. In the present investigation we exposed male and female Fischer 344 rats in a transverse electromagnetic transmission line chamber to microwaves of 915 MHz as continuous wave (CW) and pulse-modulated with repetition rates of 8, 16, 50, and 200 s⁻¹. The specific energy absorption rate (SAR) varied between 0.016 and 5 W/kg. The rats were not anesthetized during the 2-hour exposure. All animals were sacrificed by perfusion-fixation of the brains under chloral hydrate anesthesia about 1 hour after the exposure. The brains were perfused with saline for 3-4 minutes, and thereafter fixed in 4% formaldehyde for 5-6 minutes. Central coronal sections of the brains were dehydrated and embedded in paraffin and sectioned at 5 microns. Albumin and fibrinogen were demonstrated immunohistochemically. The results show albumin leakage in 5 of 62 of the controls and in 56 of 184 of the animals exposed to 915 MHz microwaves. Continuous wave resulted in 14 positive findings of 35, which differ significantly from the controls (P = 0.002).

Salford LG, Brun A, Persson BRR, Brain tumour development in rats exposed to electromagnetic fields used in wireless cellular communication. *Wireless network* 3: 463-469, 1997.

It has been suggested that electromagnetic fields (EMF) act as promoters late in the carcinogenesis process. To date, however, there is no convincing laboratory evidence that EMFs cause tumour promotion at non-thermal exposure levels. Therefore the effects of exposure to electromagnetic fields were investigated in a rat brain glioma model. Some of the exposures correspond to electromagnetic fields used in wireless communication. Microwaves at 915 MHz were used both as continuous waves (1 W), and pulse-modulated at 4, 8, 16 and 217 Hz in 0.57 ms pulses and 50 Hz in 6.67 ms pulses (2 W per pulse). Fischer 344 rats of both sexes were used in the experiments. By stereotaxic technique rat glioma cells (RG2 and N32) were injected into the head of the right caudate nucleus in 154 pairs of rats, exposed and matched controls. Starting on day 5 after inoculation, the animals were exposed for 7 hours a day, 5 days a week during 2-3 weeks. Exposed animals were kept unanaesthetized in well-ventilated TEM cells producing 915 MHz continuous or modulated microwaves. Their matched controls were kept in identical TEM cells without EMF exposure. All brains were examined histopathologically and the tumour size was estimated as the volume of an ellipsoid. Our study of 154 matched pairs of rats does not show any significant difference in tumour size between animals exposed to 915 MHz, and those not exposed. Thus our results do not support that even an extensive daily exposure to EMF promotes tumour growth when given from the fifth day after the start of tumour growth in the rat brain until the sacrifice of the animal after about 16 days.

Salford LG, Brun AR, Eberhardt JL, Malmgren L, Persson BRR, Nerve cell damage in mammalian brain after exposure to microwaves from GSM mobile phones. *Environ Health Persp* 111:881-883, 2003.

The possible risks of radio-frequency electromagnetic fields for the human body is a growing concern for the society. We have earlier shown that weak pulsed microwaves give rise to a significant leakage of albumin through the blood-brain barrier (BBB). Now we have investigated whether a pathological leakage over the BBB might be combined with damage to the neurons. Three groups of each 8 rats were exposed for 2 hours to GSM mobile phone electromagnetic fields of different strengths. We found, and present here for the first time, highly significant ($p < 0.002$) evidence for neuronal damage in both the cortex, the hippocampus and the basal ganglia in the brains of exposed rats.

Sandstrom M, Wilen J, Oftedal G, Hansson Mild K, Mobile phone use and subjective symptoms. Comparison of symptoms experienced by users of analogue and digital mobile phones. *Occup Med (Lond)* 51(1):25-35, 2001.

In 1995 many people reported symptoms such as headaches, feelings of discomfort, warmth behind/around or on the ear and difficulties concentrating while using mobile phones. The number of complaints was higher for people using the digital (GSM) system, i.e. with pulse modulated fields, than for those using the analogue (NMT) system. Our main hypothesis was that GSM users

experience more symptoms than NMT users. An epidemiological investigation was initiated including 6379 GSM users and 5613 NMT 900 users in Sweden, and 2500 from each category in Norway. The adjusted odds ratio did not indicate any increased risk for symptoms for GSM users compared with NMT 900 users. Our hypothesis was therefore disproved. However, we observed a statistically significant lower risk for sensations of warmth on the ear for GSM users compared with NMT 900 users. The same trend was seen in Norway for sensations of warmth behind/around the ear and in Sweden for headaches and fatigue. Factors distinguishing the two systems (radio frequency emission, phone temperatures and various ergonomic factors) may be responsible for these results, as well as for a secondary finding: a statistically significant association between calling time/number of calls per day and the prevalence of warmth behind/around or on the ear, headaches and fatigue.

Santini R, Seigne M, Bonhomme-Faivre L, Bouffet S, Defrasne E, Sage M. Symptoms experienced by users of digital cellular phones: a pilot study in a French engineering school. *Pathol Biol (Paris)* 49(3):222-226, 2001.

[Article in French] A survey study, using questionnaire, was conducted in 161 students and workers in a French engineering school on symptoms experienced during use of digital cellular phones. A significant increase in concentration difficult ($p < 0.05$) was reported by users of 1800-MHz (DCS) cellular phones compared to 900-MHz (GSM) phone users. In users of cellular phones, women significantly ($p < 0.05$) complained more often of sleep disturbance than men. This sex difference for sleep complaint is not observed between women and men non-users of cellular phone. The use of both cellular phones and VDT significantly ($p < 0.05$) increased concentration difficulty. Digital cellular phone users also significantly ($p < 0.05$) more often complained of discomfort, warmth, and picking on the ear during phone conversation in relation with calling duration per day and number of calls per day. The complaint warmth on the ear might be a signal to users for stopping the call.

Santini R, Santini P, Danze JM, Le Ruz P, Seigne M. Study of the health of people living in the vicinity of mobile phone base stations: I. Influence of distance and sex. *Pathol Biol (Paris)* 50(6):369-373, 2002. [Article in French]

A survey study using questionnaire was conducted in 530 people (270 men, 260 women) living or not in vicinity of cellular phone base stations, on 18 Non Specific Health Symptoms. Comparisons of complaints frequencies (CHI-SQUARE test with Yates correction) in relation with distance from base station and sex, show significant ($p < 0.05$) increase as compared to people living > 300 m or not exposed to base station, till 300 m for tiredness, 200 m for headache, sleep disturbance, discomfort, etc. 100 m for irritability, depression, loss of memory, dizziness, libido decrease, etc. Women significantly more often than men ($p < 0.05$) complained of headache, nausea, loss of appetite, sleep disturbance, depression, discomfort and visual perturbations. This first study on symptoms experienced by people living in vicinity of base stations shows that, in view of radioprotection, minimal distance of people from cellular phone base stations should not be < 300 m.

Schirmacher A, Winters S, Fischer S, Goeke J, Galla H, Kullnick U, Ringelstein EB, Stogbauer F, Electromagnetic fields (1.8 GHz) increase the permeability to sucrose of the blood-brain barrier in vitro. *Bioelectromagnetics* 21(5):338-345, 2000.

We report an investigation on the influence of high frequency electromagnetic fields (EMF) on the permeability of an in vitro model of the blood-brain barrier (BBB). Our model was a co-culture consisting of rat astrocytes and porcine brain capillary endothelial cells (BCEC). Samples were characterized morphologically by scanning electron microscopy and immunocytochemistry. The BBB phenotype of the BCEC was shown by the presence of zona occludens protein (ZO-1) as a marker for tight junctions and the close contact of the cells together with the absence of intercellular clefts. Permeability measurements using (14)C-sucrose indicated a physiological tightness which correlated with the morphological findings and verified the usefulness of our in vitro model. Samples were exposed to EMF conforming to the GSM1800-standard used in mobile telephones (1.8 GHz). The permeability of the samples was monitored over four days and compared with results of samples that were cultured identically but not exposed to EMF. Exposure to EMF increased permeability for (14)C-sucrose significantly compared to unexposed samples. The underlying pathophysiological mechanism remains to be investigated.

Shallom JM, Di Carlo AL, Ko D, Penafiel LM, Nakai A, Litovitz TA. Microwave exposure induces Hsp70 and confers protection against hypoxia in chick embryos. *J Cell Biochem* 86(3):490-496, 2002. To determine if microwave exposure could elicit a biological effect in the absence of thermal stress, studies were designed in which chick embryos were exposed to athermal microwave radiation (915 MHz) to look for induction of Hsp70, a protein produced during times of cellular stress that aids in the protection of cellular components. Levels of Hsp70 were found to increase within 2 h, with maximum expression (approximately 30% higher than controls) typically occurring by 3 h from the start of exposure. Other embryos were exposed to microwave radiation prior to being subjected to hypoxic stress, and were found to have significantly higher survival ($P < 0.05$) following re-oxygenation than non-exposed controls. The results of these studies indicate that not only can athermal microwave exposures activate the stress protein response pathway; they can also enhance survivability following exposure to a subsequent, potentially lethal stress. From a public health standpoint, it is important that more studies be performed to determine if repeated exposures, a condition likely to be found in cell phone use, are still beneficial.

Sienkiewicz ZJ, Blackwell RP, Haylock RG, Saunders RD, Cobb BL, Low-level exposure to pulsed 900 MHz microwave radiation does not cause deficits in the performance of a spatial learning task in mice. *Bioelectromagnetics* 21(3):151-158, 2000.

There is some concern that short-term memory loss or other cognitive effects may be associated with the use of mobile cellular telephones. In this experiment, the effect of repeated, acute exposure to a low intensity 900 MHz radiofrequency (RF) field pulsed at 217 Hz was explored using an appetitively-motivated spatial learning and working memory task. Adult male C57BL/6J mice were exposed under far field conditions in a GTEM cell for 45 min each day for 10 days at an average whole-body specific energy absorption rate (SAR) of 0.05 W/kg. Their performance in an 8-arm radial maze was compared to that of sham-exposed control animals. All behavioral assessments were performed without handlers having knowledge of the exposure status of the animals. Animals were tested in the maze immediately following exposure or after a delay of 15 or 30 min. No significant field-dependent effects on performance were observed in choice accuracy or in total times to complete the task across the experiment. These results suggest that exposure to RF radiation simulating a digital wireless telephone (GSM) signal under the conditions of this experiment does not affect the acquisition of the learned response. Further studies are planned to explore the effects of other SARs on learned behavior.

Singh B, Bate LA, Responses of pulmonary intravascular macrophages to 915-MHz microwave radiation: ultrastructural and cytochemical study. *Anat Rec* 246(3):343-355, 1996. BACKGROUND: Microwave (MW) radiation is being increasingly used as a source of heat supplementation during early postnatal development of pigs. Although MW radiation does not cause deleterious physiological effects, no specific information exists regarding its impact on immune cells such as macrophages. Pulmonary intravascular macrophages (PIMs) are emerging as important inflammatory cells due to their endocytic and secretory potential. An in vivo study was conducted to evaluate the effects of infrared, and low and high power MW radiation on the PIMs of pigs. METHODS: Pigs were exposed to infrared (IR), low MW (LMW; 6.1mW cm⁻²), and high MW (HMW; 11.4mW cm⁻²) radiation at 915 MHz (n = 2 for each treatment) for 24 hr. The controls (n = 2) were exposed to natural light for the same period of time. Lung tissues were processed for ultrastructural examination and acid phosphatase (AcPase) cytochemistry. In addition, rough endoplasmic reticulum (RER) as a fraction of cytosol of the PIMs was counted. RESULTS: Ultrastructural and numerical data suggested enhanced secretory activity in the PIMs of LMW-treated pigs as indicated by the increased RER:cytoplasm ratio, prominent Golgi complex profiles, and accumulation of secretory vesicles in conjunction with microtubules as compared with the control, IR, and HMW-exposed pigs. High MW treatment induced some damage to pulmonary interstitium as deduced from the presence of extracellular AcPase precipitates and disrupted collagen matrix. Intracellular globules were noticed in the PIMs of IR and LMW-treated pigs but not in the control and HMW-radiated animals. CONCLUSIONS: Elaboration of structural signs of secretory activity in the PIMs by LMW radiation in the absence of pulmonary pathological changes indicates its potential for cell activation in addition to the already established role of LMW in

heat supplementation. This activation could be due to either increased core body temperature or initiation of intracellular signaling by the LMW radiation. This study also shows that the HMW radiation is capable of inducing pathology in the form of changes in the pulmonary interstitial matrix and may not be a good source of supplementary heat.

Smythe JW, Costall B. Mobile phone use facilitates memory in male, but not female, subjects. *Neuroreport* 14(2):243-246, 2003. In the present study we report on the effects of mobile phone exposure on short- and long-term memory in male and female subjects. Subjects were university undergraduate students, and consisted of right-handed, males (= 33) and females (= 29). Individuals were randomly assigned to one of three experimental conditions: no phone exposure; inactive phone exposure; and active phone exposure. They were provided with a series of words to learn, structured in a two-dimensional shape, and given 3 min to memorise the words. After a 12 min distraction task, they were then asked to draw the shape (spatial) and place the correct words (semantic) into the appropriate boxes. One week later the same subjects were brought back to again redraw the shape and words. Error scores were determined and analysed by non-parametric techniques. The results show that males exposed to an active phone made fewer spatial errors than those exposed to an active phone condition, while females were largely unaffected. These results further indicate that mobile phone exposure has functional consequences for human subjects, and these effects appear to be sex-dependent.

Stagg RB, Thomas WJ, Jones RA, Adey WR, DNA synthesis and cell proliferation in C6 glioma and primary glial cells exposed to a 836.55 MHz modulated radiofrequency field. *Bioelectromagnetics* 18(3):230-236, 1997.

We have tested the hypothesis that modulated radiofrequency (RF) fields may act as a tumor-promoting agent by altering DNA synthesis, leading to increased cell proliferation. In vitro tissue cultures of transformed and normal rat glial cells were exposed to an 836.55 MHz, packet-modulated RF field at three power densities: 0.09, 0.9, and 9 mW/cm², resulting in specific absorption rates (SARs) ranging from 0.15 to 59 μ W/g. TEM-mode transmission-line cells were powered by a prototype time-domain multiple-access (TDMA) transmitter that conforms to the North American digital cellular telephone standard. One sham and one energized TEM cell were placed in standard incubators maintained at 37 degrees C and 5% CO₂. DNA synthesis experiments at **0.59-59 μ W/g SAR** were performed on log-phase and serum-starved semiquiescent cultures after 24 h exposure. Cell growth at 0.15-15 μ W/g SAR was determined by cell counts of log-phase cultures on days 0, 1, 5, 7, 9, 12, and 14 of a 2 week protocol. Results from the DNA synthesis assays differed for the two cell types. Sham-exposed and RF-exposed cultures of primary rat glial cells showed no significant differences for either log-phase or serum-starved condition. ***C6 glioma cells exposed to RF at 5.9 μ W/g SAR (0.9 mW/cm²) exhibited small (20-40%) significant increases in 38% of [3H]thymidine incorporation experiments.*** Growth curves of sham and RF-exposed cultures showed no differences in either normal or transformed glial cells at any of the power densities tested. Cell

doubling times of C6 glioma cells [sham (21.9 \pm 1.4 h) vs. field (22.7 \pm 3.2 h)] also demonstrated no significant differences that could be attributed to altered DNA synthesis rates. Under these conditions, this modulated RF field did not increase cell proliferation of normal or transformed cultures of glial origin.

Stagg RB, Hawel LH III, Pastorian K, Cain C, Adey WR, Byus CV, Effect of Immobilization and Concurrent Exposure to a Pulse-Modulated Microwave Field on Core Body Temperature, Plasma ACTH and Corticosteroid, and Brain Ornithine Decarboxylase, Fos and Jun mRNA. *Radiat Res* 155(4):584-592, 2001. Effect of Immobilization and Concurrent Exposure to a Pulse-Modulated Microwave Field on Core Body Temperature, Plasma ACTH and Corticosteroid, and Brain Ornithine Decarboxylase, Fos and Jun mRNA. Exposure of humans and rodents to radiofrequency (RF) cell phone fields has been reported to alter a number of stress-related parameters. To study this potential relationship in more detail, tube-restrained immobilized Fischer 344 rats were exposed in the near field in a dose-dependent manner to pulse-modulated (11 packets/s) digital cell phone microwave fields at 1.6 GHz in accordance with the Iridium protocol. Core body temperatures, plasma levels of the stress-induced hormones adrenocorticotrophic hormone (ACTH) and corticosterone, and brain levels of ornithine decarboxylase (Odc), Fos and Jun mRNAs were measured as potential markers of stress responses mediated by RF radiation. We tested the effects of the loose-tube immobilization with and without prior conditioning throughout a 2-h period (required for near-field head exposure to RF fields), on core body temperature, plasma ACTH and corticosteroids. Core body temperature increased transiently (\pm 0.3 degrees C) during the initial 30 min of loose-tube restraint in conditioned animals. When conditioned/tube-trained animals were followed as a function of time after immobilization, both the ACTH and corticosterone levels were increased by nearly 10-fold. For example, within 2-3 min, ACTH increased to 83.2 \pm 31.0 pg/dl, compared to 28.1 \pm 7.7 pg/dl for cage controls, reaching a maximum at 15-30 min (254.6 \pm 46.8 pg/dl) before returning to near resting levels by 120 min (31.2 \pm 10.2 pg/dl). However, when non-tube-trained animals were submitted to loose-tube immobilization, these animals demonstrated significantly higher (3-10-fold greater) hormone levels at 120 min than their tube-trained counterparts (313.5 \pm 54.8 compared to 31.2 \pm 10.2 pg/dl; corticosterone, 12.2 \pm 6.2 μ g/dl compared to 37.1 \pm 6.4 μ g/dl). Hormone levels in exposed animals were also compared to those in swim-stressed animals. Swimming stress also resulted in marked elevation in both ACTH and corticosterone levels, which were 10-20 fold higher (541.8 compared to 27.2-59.1 pg/dl for ACTH) and 2-5 fold higher (45.7 compared to 8.4- 20.0 μ g/dl for corticosteroids) than the cage control animals. Three time-averaged brain SAR levels of 0.16, 1.6 and 5 W/ kg were tested in a single 2-h RF-field exposure to the Iridium cell phone field. When RF-exposed and sham-exposed (immobilized) animals were compared, no differences were seen in core body temperature, corticosterone or ACTH that could be attributed to near-field RF radiation. Levels of Odc, Fos and Jun mRNA were also monitored in brains of animals exposed to the RF field for 2 h, and they showed no differences from

sham-exposed (loose-tube immobilized) animals that were due to RF-field exposure. These data suggest that a significant stress response, indicated by a transient increase in core body temperature, ACTH and corticosterone, occurred in animals placed in even the mild loose-tube immobilization required for near-field RF exposure employed here and in our other studies. Failure to adequately characterize and control this immobilization response with appropriate cage control animals, as described previously, could significantly mask any potential effects mediated by the RF field on these and other stress-related parameters. We conclude that the pulse-modulated digital Iridium RF field at SARs up to 5 W/kg is incapable of altering these stress-related responses. This conclusion is further supported by our use of an RF-field exposure apparatus that minimized immobilization stress; the use of conditioned/tube-trained animals and the measurement of hormonal and molecular markers after 2 h RF-field exposure when the stress-mediated effects were complete further support our conclusion.

Stang A, Anastassiou G, Ahrens W, Broman K, Bornfeld N, Jockel KH, The possible role of radiofrequency radiation in the development of uveal melanoma. *Epidemiology* 12(1):7-12, 2001. There are few epidemiologic studies dealing with electromagnetic radiation and uveal melanoma. The majority of these studies are exploratory and are based on job and industry titles only. We conducted a hospital-based and population-based case-control study of uveal melanoma and occupational exposures to different sources of electromagnetic radiation, including radiofrequency radiation. We then pooled these results. We interviewed a total of 118 female and male cases with uveal melanoma and 475 controls matching on sex, age, and study regions. Exposure to radiofrequency-transmitting devices was rated as (a) no radiofrequency radiation exposure, (b) possible exposure to mobile phones, or (c) probable/certain exposure to mobile phones. Exposures were rated independently by two of the authors who did not know case or control status. We used conditional logistic regression to calculate odds ratios (ORs) and 95% confidence intervals (95% CIs). We found an elevated risk for exposure to radiofrequency-transmitting devices (exposure to radio sets, OR = 3.0, 95% CI = 1.4-6.3; probable/certain exposure to mobile phones, OR = 4.2, 95% CI = 1.2-14.5). Other sources of electromagnetic radiation such as high-voltage lines, electrical machines, complex electrical environments, visual display terminals, or radar units were not associated with uveal melanoma. This is the first study describing an association between radiofrequency radiation exposure and uveal melanoma. Several methodologic limitations prevent our results from providing clear evidence on the hypothesized association.

Sykes PJ, McCallum BD, Bangay MJ, Hooker AM, Morley AA. Effect of Exposure to 900 MHz Radiofrequency Radiation on Intrachromosomal Recombination in pKZ1 Mice. *Radiat Res* 156(5):495-502, 2001. Radiofrequency (RF) radiation emitted from mobile phones is not considered to be directly genotoxic, but it may have downstream effects on cellular DNA. We studied the effect of 4 W/kg pulsed 900 MHz RF radiation on somatic

intrachromosomal recombination in the spleen in the pKZ1 recombination mutagenesis model. Somatic intrachromosomal recombination inversion events were detected in spleen tissue of pKZ1 mice by histochemical staining for E. coli beta-galactosidase protein in cells in which the lacZ transgene has undergone an inversion event. pKZ1 mice were exposed daily for 30 min to plane-wave fields of 900 MHz with a pulse repetition frequency of 217 Hz and a pulse width of 0.6 ms for 1, 5 or 25 days. Three days after the last exposure, spleen sections were screened for DNA inversion events. There was no significant difference between the control and treated groups in the 1- and 5-day exposure groups, but there was a significant reduction in inversions below the spontaneous frequency in the 25-day exposure group. This observation suggests that exposure to RF radiation can lead to a perturbation in recombination frequency which may have implications for recombination repair of DNA. The biological significance of a reduction below the spontaneous frequency is not known. The number of mice in each treatment group in this study was small ($n = 10$ or $n = 20$). Therefore, repetition of this study with a larger number of animals is required to confirm these observations.

Takahashi S, Inaguma S, Cho Y-M, Imaida K, Wang J, Fujiwara O, Shirai T, Lack of Mutation Induction with Exposure to 1.5 GHz Electromagnetic Near Fields Used for Cellular Phones in Brains of Big Blue Mice. *Cancer Res* 62:1956-1960, 2002.

The possible mutagenic potential of exposure to 1.5 GHz electromagnetic near field (EMF) was investigated using brain tissues of Big Blue mice (BBM). Male BBM were locally exposed to EMF in the head region at 2.0, 0.67, and 0 W/kg specific absorption rate for 90 min/day, 5 days/week, for 4 weeks. No gliosis or degenerative lesions were histopathologically noted in brain tissues, and no obvious differences in Ki-67 labeling and apoptotic indices of glial cells were evident among the groups. There was no significant variation in the frequency of independent mutations of the *lacI* transgene in the brains. G:C to A:T transitions at CpG sites constituted the most prevalent mutations in all groups and at all time points. Deletion mutations were slightly increased in both the high and low EMF exposure groups as compared with the sham-exposed group, but the differences were not statistically significant. These findings suggest that exposure to 1.5 GHz EMF is not mutagenic to mouse brain cells and does not create any increased hazard with regard to brain tumor development.

Testylier G, Tonduli L, Malabiau R, Debouzy JC. Effects of exposure to low level radiofrequency fields on acetylcholine release in hippocampus of freely moving rats. *Bioelectromagnetics* 23:249-255, 2002.

Some central cholinergic effects have been reported in animals after acute exposure to radiofrequency electromagnetic field at low intensity. We studied acetylcholine (ACh) release in the brain of freely moving rats exposed for 1 h during the day to a 2.45 GHz continuous wave radiofrequency field (RF) (2 or 4 mW/cm²) or exposed for 1 or 14 h during the night to a 800 MHz field modulated at 32 Hz (AM 200 mW/cm²). Measurements were performed by microdialysis

using a membrane implanted through the upper CA1 region of the hippocampus. After irradiation with the 2.45 GHz RF, rats exposed at 2 mW/cm² did not show a significant modification of ACh release, whereas those exposed at 4 mW/cm² showed a significant 40% decrease in mean ACh release from hippocampus. This decrease was maximal at 5 h post exposure. Exposure to the 800 MHz RF for 1 h did not cause any significant effect, but exposure for 14 hrs induced a significant 43% decrease in ACh release during the period 11 p.m.-4 a.m. compared to control rats. In the control group we observed an increase of ACh release at the beginning of the night, which was linked to the waking period of rats. This normal increase was disturbed in rats exposed overnight to the 800 MHz RF. This work indicates that neurochemical modification of the hippocampal cholinergic system can be observed during and after an exposure to low intensity RF.

Tice RR , Hook GG , Donner M , McRee DI, Guy AW. Genotoxicity of radiofrequency signals. I. Investigation of DNA damage and micronuclei induction in cultured human blood cells. *Bioelectromagnetics* 23:113-126, 2002.

As part of a comprehensive investigation of the potential genotoxicity of radiofrequency (RF) signals emitted by cellular telephones, in vitro studies evaluated the induction of DNA and chromosomal damage in human blood leukocytes and lymphocytes, respectively. The signals were voice modulated 837 MHz produced by an analog signal generator or by a time division multiple access (TDMA) cellular telephone, 837 MHz generated by a code division multiple access (CDMA) cellular telephone (not voice modulated), and voice modulated 1909.8 MHz generated by a global system of mobile communication (GSM)-type personal communication systems (PCS) cellular telephone. DNA damage (strand breaks/alkali labile sites) was assessed in leukocytes using the alkaline (pH>13) single cell gel electrophoresis (SCG) assay. Chromosomal damage was evaluated in lymphocytes mitotically stimulated to divide postexposure using the cytochalasin B-binucleate cell micronucleus assay. Cells were exposed at 37±1°C, for 3 or 24 h at average specific absorption rates (SARs) of 1.0-10.0 W/kg. Exposure for either 3 or 24 h did not induce a significant increase in DNA damage in leukocytes, nor did exposure for 3 h induce a significant increase in micronucleated cells among lymphocytes. However, exposure to each of the four RF signal technologies for 24 h at an average SAR of 5.0 or 10.0 W/kg resulted in a significant and reproducible increase in the frequency of micronucleated lymphocytes. The magnitude of the response (approximately four fold) was independent of the technology, the presence or absence of voice modulation, and the frequency (837 vs. 1909.8 MHz). This research demonstrates that, under extended exposure conditions, RF signals at an average SAR of at least 5.0 W/kg are capable of inducing chromosomal damage in human lymphocytes.

Tsurita G, Nagawa H, Ueno S, Watanabe S, Taki M, Biological and morphological effects on the brain after exposure of rats to a 1439 MHz TDMA field. *Bioelectromagnetics* 21(5):364-371, 2000.

We investigated the effects of exposure to a 1439 MHz TDMA (Time Division Multiple Access) field, as used in cellular phones, on the permeability of the blood-brain barrier (BBB), on the morphological changes of the brain, and on body-mass fluctuations. Male Sprague-Dawley (SD) rats were divided into Three groups of eight rats each. The rats in the EM(+) group, which had their heads arrayed in a circle near the central antenna of an exposure system, were exposed to a 1439 MHz field for one hour a day. The rats in EM(-) group were also in the exposure system, however, without high-frequency electromagnetic wave (HF-EMW) exposure. The animals in the control group were neither placed in the system nor exposed to HF-EMWs. The exposure period was two or four weeks. The energy dose rate peaked at 2 W/kg in the brain; the average over the whole body was 0.25 W/kg. The changes in the permeability of BBB were investigated by Evans blue injection method and by immunostaining of serum albumin. HF-EMWs had no effect on the permeability of BBB. The morphological changes in the cerebellum were investigated by assessing the degeneration of Purkinje cells and the cell concentration in the granular layer. No significant changes were observed in the groups of rats exposed to HF-EMWs for two or four weeks. Averaged body masses were not affected by HF-EMWs exposure. In conclusion, a 1439 MHz TDMA field did not induce observable changes in the permeability of the BBB, morphological changes in the cerebellums, or body mass changes in rats, as evaluated by the conventional methods.

Urban, P, Lukas, E, Roth, Z, Does acute exposure to the electromagnetic field emitted by a mobile phone influence visual evoked potentials? A pilot study. *Cent Eur J Public Health* 6(4):288-290, 1998.

To search for a potential negative influence on the central nervous system (CNS) of the electromagnetic field emitted by a mobile phone, the authors performed a pilot experimental study of the influence of a single short acute exposure to the GSM mobile phone Motorola 8700, using visual evoked potentials (VEP) examination as an electrophysiological marker of CNS dysfunction. The study group consisted of 20 healthy volunteers. The duration of exposure was 5 minutes. The output power of the device was 1.5 W when the antenna was pulled up. Five parameters of VEP were evaluated by means of multifactorial ANOVA. Confounding effects of age, sex, and of the call in itself were taken into consideration. No statistically significant influence of the above-described exposure to the electromagnetic field emitted by the mobile phone on latencies or amplitudes of VEP was observed.

Utteridge TD, Gebiski V, Finnie JW, Vernon-Roberts B, Kuchel TR. Long-Term Exposure of E_Pim1 Transgenic Mice to 898.4 MHz Microwaves does not Increase Lymphoma Incidence. *Radiat Res* 158(3):357-364, 2002.

A total of 120 E₋Pim1 heterozygous mice and 120 wild-type mice were exposed for 1 h/day 5 days/week at each of the four exposure levels in "Ferris-wheel" exposure systems for up to 104 weeks to GSM-modulated 898.4 MHz radiation at SARs of 0.25, 1.0, 2.0 and 4.0 W/kg. In addition, 120 heterozygous and 120 wild-type mice were sham-exposed; there was also an unrestrained negative control group. Four exposure levels were used to investigate whether a dose-response effect could be detected. Independent verification confirmed that the exposures in the current study were nonthermal. There was no significant difference in the incidence of lymphomas between exposed and sham-exposed groups at any of the exposure levels. A dose-response effect was not detected. The findings showed that long-term exposures of lymphoma-prone mice to 898.4 MHz GSM radiofrequency (RF) radiation at SARs of 0.25, 1.0, 2.0 and 4.0 W/kg had no significant effects when compared to sham-irradiated animals. A previous study (Repacholi et al., Radiat. Res. 147, 631-640, 1997) reported that long-term exposure of lymphoma-prone mice to one exposure level of 900 MHz RF radiation significantly increased the incidence of non-lymphoblastic lymphomas when compared to sham-irradiated animals.

Velizarov, S, Raskmark, P, Kwee, S, The effects of radiofrequency fields on cell proliferation are non-thermal. *Bioelectrochem Bioenerg* 48(1):177-180, 1999.

The number of reports on the effects induced by radiofrequency (RF) electromagnetic fields and microwave (MW) radiation in various cellular systems is still increasing. Until now no satisfactory mechanism has been proposed to explain the biological effects of these fields. One of the current theories is that heat generation by RF/MW is the cause, in spite of the fact that a great number of studies under isothermal conditions have reported significant cellular changes after exposure to RF/MW. Therefore, this study was undertaken to investigate which effect MW radiation from these fields in combination with a significant change of temperature could have on cell proliferation. The experiments were performed on the same cell line, and with the same exposure system as in a previous work [S. Kwee, P. Raskmark, Changes in cell proliferation due to environmental non-ionizing radiation: 2. Microwave radiation, *Bioelectrochem. Bioenerg.*, 44 (1998), pp. 251-255]. The field was generated by signal simulation of the Global System for Mobile communications (GSM) of 960 MHz. Cell cultures, growing in microtiter plates, were exposed in a specially constructed chamber, a Transverse Electromagnetic (TEM) cell. The Specific Absorption Rate (SAR) value for each cell well was calculated for this exposure system. However, in this study the cells were exposed to the field at a higher or lower temperature than the temperature in the field-free incubator i.e., the temperature in the TEM cell was either 39 or 35 +/- 0.1 degrees C. The corresponding sham experiments were performed under exactly the same experimental conditions. The results showed that there was a significant change in cell proliferation in the exposed cells in comparison to the non-exposed (control) cells at both

temperatures. On the other hand, no significant change in proliferation rate was found in the sham-exposed cells at both temperatures. This shows that biological effects due to RF/MW cannot be attributed only to a change of temperature. Since the RF/MW induced changes were of the same order of magnitude at both temperatures and also comparable to our previous results under isothermal conditions at 37 degrees C, cellular stress caused by electromagnetic fields could initiate the changes in cell cycle reaction rates. It is widely accepted that certain classes of heat-shock proteins are involved in these stress reactions.

Von Klitzing, L, Low-frequency pulsed electromagnetic fields influence EEG of man. *Phys. Medica* 11:77-80, 1995.

New techniques using low-frequency pulsed electromagnetic fields (e.g., digital telecommunication) have raised the question for interference with the biological system of man. EEG data of man sampled under the influence of these electromagnetic fields are altered extremely in the range of alpha-activity as well as during after exposure for some hours. The biological effect is induced by field intensities lower than the given international limiting values.

Vijayalaxmi , Leal BZ, Meltz ML, Pickard WF, Bisht KS, Roti Roti JL , Straube WL, Moros EG, Cytogenetic Studies in Human Blood Lymphocytes Exposed In Vitro to Radiofrequency Radiation at a Cellular Telephone Frequency (835.62 MHz, FDMA). *Radiat Res* 155(1):113-121, 2001.

Vijayalaxmi, Pickard, W. F., Bisht, K. S., Leal, B. Z., Meltz, M. L., Roti Roti, J. L., Straube, W. L. and Moros, E. G. Cytogenetic Studies in Human Blood Lymphocytes Exposed In Vitro to Radiofrequency Radiation at a Cellular Telephone Frequency (835.62 MHz, FDMA). Freshly collected peripheral blood samples from four healthy human volunteers were diluted with RPMI 1640 tissue culture medium and exposed in sterile T-75 tissue culture flasks in vitro for 24 h to 835.62 MHz radiofrequency (RF) radiation, a frequency employed for customer-to-base station transmission of cellular telephone communications. An analog signal was used, and the access technology was frequency division multiple access (FDMA, continuous wave). A nominal net forward power of 68 W was used, and the nominal power density at the center of the exposure flask was 860 W/m². The mean specific absorption rate in the exposure flask was 4.4 or 5.0 W/kg. Aliquots of diluted blood that were sham-exposed or exposed in vitro to an acute dose of 1.50 Gy of gamma radiation were used as negative or positive controls. Immediately after the exposures, the lymphocytes were stimulated with a mitogen, phytohemagglutinin, and cultured for 48 or 72 h to determine the extent of genetic damage, as assessed from the frequencies of chromosomal aberrations and micronuclei. The extent of alteration in the kinetics of cell proliferation was determined from the mitotic indices in 48-h cultures and from the incidence of binucleate cells in 72-h cultures. The data indicated no significant differences between RF-radiation- and sham-exposed lymphocytes with respect to mitotic indices, incidence of exchange aberrations, excess fragments, binucleate cells, and micronuclei. In contrast, the response of the

lymphocytes exposed to gamma radiation was significantly different from both RF-radiation- and sham-exposed cells for all of these indices. Thus, under the experimental conditions tested, there is no evidence for the induction of chromosomal aberrations and micronuclei in human blood lymphocytes exposed in vitro for 24 h to 835.62 MHz RF radiation at SARs of 4.4 or 5.0 W/kg.

Vijayalaxmi, Bisht KS, Pickard WF, Meltz ML, Roti Roti JL, Moros EG. Chromosome damage and micronucleus formation in human blood lymphocytes exposed in vitro to radiofrequency radiation at a cellular telephone frequency (847.74 MHz, CDMA). *Radiat Res* 156(4):430-432, 2001.

Peripheral blood samples collected from four healthy nonsmoking human volunteers were diluted with tissue culture medium and exposed in vitro for 24 h to 847.74 MHz radiofrequency (RF) radiation (continuous wave), a frequency employed for cellular telephone communications. A code division multiple access (CDMA) technology was used with a nominal net forward power of 75 W and a nominal power density of 950 W/m² (95 mW/cm²). The mean specific absorption rate (SAR) was 4.9 or 5.5 W/kg. Blood aliquots that were sham-exposed or exposed in vitro to an acute dose of 1.5 Gy of gamma radiation were included in the study as controls. The temperatures of the medium during RF-radiation and sham exposures in the Radial Transmission Line facility were controlled at 37 +/- 0.3 degrees C. Immediately after the exposures, lymphocytes were cultured at 37 +/- 1 degrees C for 48 or 72 h. The extent of genetic damage was assessed from the incidence of chromosome aberrations and micronuclei. The kinetics of cell proliferation was determined from the mitotic indices in 48-h cultures and from the incidence of binucleate cells in 72-h cultures. The data indicated no significant differences between RF-radiation-exposed and sham-exposed lymphocytes with respect to mitotic indices, frequencies of exchange aberrations, excess fragments, binucleate cells, and micronuclei. The response of gamma-irradiated lymphocytes was significantly different from that of both RF-radiation-exposed and sham-exposed cells for all of these indices. Thus there was no evidence for induction of chromosome aberrations and micronuclei in human blood lymphocytes exposed in vitro for 24 h to 847.74 MHz RF radiation (CDMA) at SARs of 4.9 or 5.5 W/kg.

Vijayalaxmi, Sasser LB, Morris JE, Wilson BW, Anderson LE. Genotoxic Potential of 1.6 GHz Wireless Communication Signal: In Vivo Two-Year Bioassay. *Radiat Res* 159(4):558-564, 2003.

Timed-pregnant Fischer 344 rats (from nineteenth day of gestation) and their nursing offspring (until weaning) were exposed to a far-field 1.6 GHz Iridium wireless communication signal for 2 h/day, 7 days/week. Far-field whole-body exposures were conducted with a field intensity of 0.43 mW/cm² and whole-body average specific absorption rate (SAR) of 0.036 to 0.077 W/kg (0.10 to 0.22 W/kg in the brain). This was followed by chronic, head-only exposures of male and female offspring to a near-field 1.6 GHz signal for 2 h/day, 5 days/week, over 2 years. Near-field exposures were conducted at an SAR of 0.16 or 1.6 W/kg in the brain. Concurrent sham-exposed and cage control rats were also included in

the study. At the end of 2 years, all rats were necropsied. Bone marrow smears were examined for the extent of genotoxicity, assessed from the presence of micronuclei in polychromatic erythrocytes. The results indicated that the incidence of micronuclei/2000 polychromatic erythrocytes were not significantly different between 1.6 GHz-exposed, sham-exposed and cage control rats. The group mean frequencies were 5.6 +/- 1.8 (130 rats exposed to 1.6 GHz at 0.16 W/kg SAR), 5.4 +/- 1.5 (135 rats exposed to 1.6 GHz at 1.6 W/kg SAR), 5.6 +/- 1.7 (119 sham-exposed rats), and 5.8 +/- 1.8 (100 cage control rats). In contrast, positive control rats treated with mitomycin C exhibited significantly elevated incidence of micronuclei/2000 polychromatic erythrocytes in bone marrow cells; the mean frequency was 38.2 +/- 7.0 (five rats). Thus there was no evidence for excess genotoxicity in rats that were chronically exposed to 1.6 GHz compared to sham-exposed and cage controls.

Vollrath L, Spessert R, Kratzsch T, Keiner M, Hollmann H, No short-term effects of high-frequency electromagnetic fields on the mammalian pineal gland. *Bioelectromagnetics* 18(5):376-387, 1997.

There is ample experimental evidence that changes of earth-strength static magnetic fields, pulsed magnetic fields, or alternating electric fields (60 Hz) depress the nocturnally enhanced melatonin synthesis of the pineal gland of certain mammals. No data on the effects of high-frequency electromagnetic fields on melatonin synthesis is available. In the present study, exposure to 900 MHz electromagnetic fields [0.1 to 0.6 mW/cm², approximately 0.06 to 0.36 W/kg specific absorption rate (SAR) in rats and 0.04 W/kg in Djungarian hamsters; both continuous and/or pulsed at 217 Hz, for 15 min to 6 hr at day or night had no notable short-term effect on pineal melatonin synthesis in male and female Sprague-Dawley rats and Djungarian hamsters. Pineal synaptic ribbon

profile numbers (studied in rats only) were likewise not affected. The 900 MHz electromagnetic fields, unpulsed or pulsed at 217 Hz, as applied in the present study, have no short-term effect on the mammalian pineal gland.

Wagner, P, Roschke, J, Mann, K, Hiller, W, Frank, C, Human sleep under the influence of pulsed radiofrequency electromagnetic fields: a polysomnographic study using standardized conditions. *Bioelectromagnetics* 19(3):199-202, 1998.

To investigate the influence of radiofrequency electromagnetic fields (EMFs) of cellular phone GSM signals on human sleep electroencephalographic (EEG) pattern, all-night polysomnographies of 24 healthy male subjects were recorded, both with and without exposure to a circular polarized EMF (900 MHz, pulsed with a frequency of 217 Hz, pulse width 577 micros, power flux density 0.2 W/m²). Suppression of rapid eye movement (REM) sleep as well as a sleep-inducing effect under field exposure did not reach statistical significance, so that previous results indicating alterations of these sleep parameters could not be replicated. Spectral power analysis also did not reveal any alterations of the EEG rhythms

during EMF exposure. The failure to confirm our previous results might be due to dose-dependent effects of the EMF on the human sleep profile.

Wagner P, Roschke J, Mann K, Fell J, Hiller W, Frank C, Grozinger M, Human sleep EEG under the influence of pulsed radio frequency electromagnetic fields. results from polysomnographies using submaximal high power flux densities. *Neuropsychobiology* 42(4):207-212, 2000. Former exploratory investigations of sleep alterations due to global system for mobile communications (GSM) signals have shown a hypnotic and REM-suppressive effect under field exposure. This effect was observed in a first study using a power flux density of 0.5 W/m², and the same trend occurred in a second study with a power flux density of 0.2 W/m². For the present study, we applied a submaximal power flux density of 50 W/m². To investigate putative effects of radio frequency electromagnetic fields (EMFs) of cellular GSM phones on human sleep EEG pattern, all-night polysomnographies of 20 healthy male subjects both with and without exposure to a circularly polarized EMF (900 MHz, pulsed with a frequency of 217 Hz, pulse duration 577 &mgr;s) were recorded. The results showed no significant effect of the field application either on conventional sleep parameters or on sleep EEG power spectra.

Warren HG, Prevatt AA, Daly KA, Antonelli PJ. Cellular telephone use and risk of intratemporal facial nerve tumor. *Laryngoscope* 113(4):663-667, 2003. (G-C, H-U, C-P) OBJECTIVES/HYPOTHESISMicrowave radiation exposure from cellular telephone use has been implicated in the development of intracranial tumors. The intratemporal facial nerve (IFN) is exposed to higher levels of cellular telephone radiation than intracranial tissues. The purpose of the study was to determine whether cellular telephone use is associated with an increased risk of IFN tumors. STUDY DESIGNCase-control using a structured telephone survey at an academic, tertiary-care referral center. METHODS Patients with IFN tumors (n = 18) were case-matched with patients treated for acoustic neuroma (n = 51), rhinosinusitis (n = 72), and dysphonia or gastroesophageal reflux disease (n = 69). Risk of facial nerve tumorigenesis was compared by extent of cellular telephone use and other risk factors.RESULTSThe odds ratio of developing an IFN tumor was 0.6 (95% CI, 0.2-1.9) with any handheld cellular telephone use and 0.4 (95% CI, 0.1-2.1) with regular cellular telephone use. No factors were associated with an increased risk for IFN tumor development. CONCLUSIONS Regular cellular telephone use does not appear to be associated with a higher risk of IFN tumor development. The short duration of widespread cellular telephone use precludes definite exclusion as a risk for IFN tumor development.

Weisbrot D, Lin H, Ye L, Blank M, Goodman R. Effects of mobile phone radiation on reproduction and development in *Drosophila melanogaster*. *J Cell Biochem* 89(1):48-55, 2003. In this report we examined the effects of a discontinuous radio frequency (RF) signal produced by a GSM multiband mobile phone (900/1,900 MHz; SAR approximately 1.4 W/kg) on *Drosophila*

melanogaster, during the 10-day developmental period from egg laying through pupation. As found earlier with low frequency exposures, the non-thermal radiation from the GSM mobile phone increased numbers of offspring, elevated hsp70 levels, increased serum response element (SRE) DNA-binding and induced the phosphorylation of the nuclear transcription factor, ELK-1. The rapid induction of hsp70 within minutes, by a non-thermal stress, together with identified components of signal transduction pathways, provide sensitive and reliable biomarkers that could serve as the basis for realistic mobile phone safety guidelines.

Wilén J, Sandström M, Hansson Mild K. Subjective symptoms among mobile phone users-A consequence of absorption of radiofrequency fields? *Bioelectromagnetics* 24(3):152-159, 2003. In a previous epidemiological study, where we studied the prevalence of subjective symptoms among mobile phone (MP) users, we found as an interesting side finding that the prevalence of many of the subjective symptoms increased with increasing calling time and number of calls per day. In this extrapolative study, we have selected 2402 people from the epidemiological study who used any of the four most common GSM MP. We used the information about the prevalence of symptoms, calling time per day, and number of calls per day and combined it with measurements of the Specific Absorption Rate (SAR). We defined three volumes in the head and measured the maximum SAR averaged over a cube of 1 g tissue (SAR(1g)) in each volume. Two new exposure parameters Specific Absorption per Day (SAD) and Specific Absorption per Call (SAC) have been devised and are obtained as combinations of SAR, calling time per day, and number of calls per day, respectively. The results indicate that SAR values >0.5 W/kg may be an important factor for the prevalence of some of the symptoms, especially in combination with long calling times per day.

Wolke S, Neibig U, Elsner R, Gollnick F, Meyer R, Calcium homeostasis of isolated heart muscle cells exposed to pulsed high-frequency electromagnetic fields. *Bioelectromagnetics* 17(2):144-153, 1996.

The intracellular calcium concentration ($[Ca^{2+}]_i$) of isolated ventricular cardiac myocytes of the guinea pig was measured during the application of pulsed high-frequency electromagnetic fields. The high-frequency fields were applied in a transverse electromagnetic cell designed to allow microscopic observation of the myocytes during the presence of the high-frequency fields. The $[Ca^{2+}]_i$ was measured as fura-2 fluorescence by means of digital image analysis. Both the carrier frequency and the square-wave pulse-modulation pattern were varied during the experiments (carrier frequencies: 900, 1,300, and 1,800 MHz pulse modulated at 217 Hz with 14 percent duty cycle; pulsation pattern at 900 MHz: continuous wave, 16 Hz, and 50 Hz modulation with 50

percent duty cycle and 30 kHz modulation with 80 percent duty cycle). The mean specific absorption rate (SAR) values in the solution were within one order of magnitude of **1 mW/kg**. They varied depending on the applied carrier frequency and pulse pattern. The experiments were designed in three phases: 500 s of sham exposure, followed by 500 s of field exposure, then chemical stimulation without field. The chemical stimulation (K⁺ -depolarization) indicated the viability of the cells. The K⁺ depolarization yielded a significant increase in [Ca(2⁺)]_i. Significant differences between sham exposure and high-frequency field exposure were not found except when a very small but statistically significant difference was detected in the case of 900 MHz/50 Hz. However, this small difference was not regarded as a relevant effect of the exposure.

Yamaguchi H, Tsurita G, Ueno S, Watanabe S, Wake K, Taki M, Nagawa H. 1439 MHz pulsed TDMA fields affect performance of rats in a T-maze task only when body temperature is elevated. *Bioelectromagnetics* 24(4):223-230, 2003. This study sought to clarify the effects of exposure to electromagnetic waves (EMW) used in cellular phones on learning and memory processes. Sprague-Dawley rats were exposed for either 1 h daily for 4 days or for 4 weeks to a pulsed 1439 MHz time division multiple access (TDMA) field in a carousel type exposure system. At the brain, average specific absorption rate (SAR) was 7.5 W/kg, and the whole body average SAR was 1.7 W/kg. Other subjects were exposed at the brain average SAR of 25 W/kg and the whole body average SAR of 5.7 W/kg for 45 min daily for 4 days. Learning and memory were evaluated by reversal learning in a food rewarded T-maze, in which rats learned the location of food (right or left) by using environmental cues. The animals exposed to EMW with the brain average SAR of 25 W/kg for 4 days showed statistically significant decreases in the transition in number of correct choices in the reversal task, compared to sham exposed or cage control animals. However, rats exposed to the brain average SAR of 7.5 W/kg for either 4 days or for 4 weeks showed no T-maze performance impairments. Intraperitoneal temperatures, as measured by a fiber optic thermometer, increased in the rats exposed to the brain average SAR of 25 W/kg but remained the same for the brain average SAR of 7.5 W/kg. The SAR of a standard cellular phone is restricted to a maximum of 2 W/kg averaged over 10 g tissue. These results suggest that the exposure to a TDMA field at levels about four times stronger than emitted by cellular phones does not affect the learning and memory processes when there are no thermal effects.

Zook BC, Simmens SJ, The Effects of 860 MHz Radiofrequency Radiation on the Induction or Promotion of Brain Tumors and Other Neoplasms in Rats. *Radiat Res* 155(4):572-583, 2001. Zook, B. C. and Simmens, S. J. The Effects of 860 MHz Radiofrequency Radiation on the Induction or Promotion of Brain Tumors and Other Neoplasms in Rats. Sprague-Dawley rats were irradiated with a continuous-wave (CW) or a pulsed-wave (P) radiofrequency (RF) for 6 h/day, 5 days/week from 2 up to 24 months of age. The RFs emanated from dipole antennas (1 W average output) 2.0 +/- 0.5 cm from the tip of each

rat's nose. The RFs had an 860 MHz frequency, and the specific absorption rate was 1.0 W/ kg averaged over the brain. Fifteen groups of 60 rats (900 total) were formed from offspring of females injected i.v. with 0 (groups 1, 2, 9, 10, 13), 2.5 (groups 5, 6, 7, 8, 11, 12, 14) or 10 mg/kg (groups 3, 4, 15) ethylnitrosourea (ENU) to induce brain tumors. Groups 1, 3, 5 and 7 received the PRF, and groups 9 and 11 the CWRF; groups 2, 4, 6, 8, 10 and 12 were sham-irradiated, and groups 13-15 were cage controls. All rats but 2, totaling 898, were necropsied, and major tissues were studied histopathologically. There was no statistically significant evidence that the PRF or CWRF induced neoplasia in any tissues. Additionally, there was no significant evidence of promotion of cranial or spinal nerve or spinal cord tumors. The PRF or CWRF had no statistically significant effect on the number, volume, location, multiplicity, histological type, malignancy or fatality of brain tumors. There was a trend for the group that received a high dose of ENU and was exposed to the PRF to develop fatal brain tumors at a higher rate than its sham group; however, the result was not significant using the log-rank test ($P = 0.14$, 2-tailed). No statistically significant differences were related to the PRF or CWRF compared to controls in the low- or zero-dose groups regarding tumors of any kind.

Dosimetry studies

Anderson V, Joyner KH, Specific absorption rate levels measured in a phantom head exposed to radio frequency transmissions from analog hand-held mobile phones. *Bioelectromagnetics* 16(1):60-69, 1995. Electric fields (E-fields) induced within a phantom head from exposure to three different advanced mobile phone system (AMPS) hand-held telephones were measured using an implantable E-field probe. Measurements were taken in the eye nearest the phone and along a lateral scan through the brain from its centre to the side nearest the phone. During measurement, the phones were positioned alongside the phantom head as in typical use and were configured to transmit at maximum power (600 mW nominal). The specific absorption rate (SAR) was calculated from the in situ E-field measurements, which varied significantly between phone models and antenna configuration. The SARs induced in the eye ranged from 0.007 to 0.21 W/kg. Metal-framed spectacles enhanced SAR levels in the eye by 9-29%. In the brain, maximum levels were recorded at the measurement point closest to the phone and ranged from 0.12 to 0.83 W/kg. These SARs are below peak spatial limits recommended in the U.S. and Australian national standards [IEEE Standards Coordinating Committee 28 (1991): C95.1-1991 and Standards Australia (1990): AS2772.1-1990] and the IRPA guidelines for safe exposure to radio frequency (RF) electromagnetic fields [IRPA (1988): Health Phys 54:115-123]. Furthermore, a detailed thermal analysis of the eye indicated only a 0.022 degrees C maximum steady-state temperature rise in the eye from a uniform SAR loading of 0.21 W/kg. A more approximate thermal analysis in the brain also indicated only a small maximum temperature rise of 0.034 degrees C for a local SAR loading of 0.83 W/kg.

Dimbylow PJ, Mann SM. SAR calculations in an anatomically realistic model of the head for mobile communication transceivers at 900 MHz and 1.8 GHz. *Phys. Med. Biol.* 39:1537-1553, 1994.

Abstract. A new mathematical model of the head has been constructed from a set of serial MRI slices from one subject. Finite-difference time-domain (FDTD) calculations of the specific energy absorption rate (SAR) have been performed on this model with a 2 mm resolution for a generic mobile communication transceiver represented by a quarter-wavelength monopole on a metal box. The antenna was mounted either at the centre or corner of the top face of the box. The frequencies considered were 900 MHz and 1.8 GHz. Three irradiation geometries were considered, a vertical handset in front of the eye and vertical and horizontal orientations at the side of the ear. The effect of a hand grasping the handset was considered. The head model was scaled to represent the head of an infant and a subset of calculations was performed to verify that the SAR deposited in the infant head did not exceed that in the adult. Results are also presented for a half-wavelength dipole. The maximum SAR values produced by the generic transceiver for the horizontal orientation at the side of the head which is the most typical position, averaged over 10 g of tissue at 900 MHz and 1.8 GHz, are 2.1 and 3.0 W kg⁻¹ per W of radiated power. The corresponding values over 1 g of tissue are 2.3 and 4.8 W kg⁻¹ per W at 900 MHz and 1.8 GHz. However, if one were to consider all possible operational conditions, the placement of the transceiver in front of the eye will give 3.1 and 4.6 W kg⁻¹ per W averaged over 10 g of tissue and 4.7 and 7.7 W kg⁻¹ per W over 1 g of tissue at 900 MHz and 1.8 GHz, respectively.

Gandhi OP, Lazzi G, Tinniswood A, Yu QS, Comparison of numerical and experimental methods for determination of SAR and radiation patterns of handheld wireless telephones. *Bioelectromagnetics Suppl* 4:93-101, 1999.

Some recent developments in both the numerical and experimental methods for determination of SARs and radiation patterns of handheld wireless telephones are described, with emphasis on comparison of results using the two methods. For numerical calculations, it was possible to use the Pro-Engineer CAD Files of cellular telephones for a realistic description of the device. Also, we used the expanding grid formulation of the finite-difference time-domain (FDTD) method for finer-resolution representation of the coupled region, including the antenna, and an increasingly coarser representation of the more-distant, less-coupled region. Together with the truncation of the model of the head, this procedure led to a saving of computer memory needed for SAR calculations by a factor of over 20. Automated SAR and radiation pattern measurement systems were used to validate both the calculated 1-g SARs and radiation patterns for several telephones, including some research test samples, using a variety of antennas. Even though widely different peak 1-g SARs were obtained, ranging from 0.13 to 5.41 W/kg, agreement between the calculated and the measured data for these telephones, five each at 835 and 1900 MHz, was excellent and generally within +/-20% (+/-1 dB). An important observation was that for a maximum radiated power of 600 mW at 800/900 MHz, which may be used for telephones using

AMPS technology, the peak 1-g SARs can be higher than 1.6 W/kg unless antennas are carefully designed and placed further away from the head.

Van de Kamer JB, Lagendijk, JJW, Computation of high-resolution SAR distributions in a head due to a radiating dipole antenna representing a hand-held mobile phone. *Phys. Med. Biol.* 47:1827-1835, 2002.

SAR distributions in a healthy female adult head as a result of a radiating vertical dipole antenna (frequency 915 MHz) representing a hand-held mobile phone have been computed for three different resolutions: 2 mm, 1 mm and 0.4 mm. The extremely high resolution of 0.4 mm was obtained with our quasistatic zooming technique, which is briefly described in this paper. For an effectively transmitted power of 0.25 W, the maximum averaged SAR values in both cubic- and arbitrary-shaped volumes are, respectively, about 1.72 and 2.55 W kg⁻¹ for 1 g and 0.98 and 1.73 W kg⁻¹ for 10 g of tissue. These numbers do not vary much (<8%) for the different resolutions, indicating that SAR computations at a resolution of 2 mm are sufficiently accurate to describe the large-scale distribution. However, considering the detailed SAR pattern in the head, large differences may occur if high-resolution computations are performed rather than low-resolution ones. These deviations are caused by both increased modelling accuracy and improved anatomical description in higher resolution simulations. For example, the SAR profile across a boundary between tissues with high dielectric contrast is much more accurately described at higher resolutions. Furthermore, low-resolution dielectric geometries may suffer from loss of anatomical detail, which greatly affects small-scale SAR distributions. Thus, for strongly inhomogeneous regions high-resolution SAR modelling is an absolute necessity.